

Twisted cohomology pairings of knots II; to classical invariants

Takefumi Nosaka

Abstract

We will show that the bilinear cup products with local coefficients of closed 3-manifolds recovers some classical invariants, including the Blanchfield pairing in the knot case and the twisted cup products of infinite covers, and the Casson-Gordon local signatures. As a result, we further give diagrammatic computations of these classical subjects.

Keywords

Cup product, knot, Blanchfield pairing, infinite covering, slice genus

Contents

1	Introduction	2
2	Result I: two relations to infinite cyclic coverings	3
2.1	Abelian case; recover of the Blanchfield pairing of knots	4
2.2	Recover of twisted cup products of infinite cyclic covers	4
3	Result II; Recover of the Casson-Gordon local signature	6
3.1	Review; the Casson-Gordon invariants and σ -signature	7
3.2	A diagrammatic formulation	8
4	Examples of computations	9
4.1	Reviews; diagrammatic computation of the twisted pairings.	9
4.2	The Blanchfield pairing of the torus knot	10
4.3	The figure eight knot	12
4.4	A computation of the Casson-Gordon local signature	12
5	Proofs of the statements in Sections 2-3	14
5.1	Proofs of the statements in Section 2.2	14
5.2	Proofs of Theorem 2.1	17
5.3	Proof of Theorem 3.1	19
A	Appendix: comparison with the work of Goldman	20
B	Universality of Alexander quandle cocycle invariants.	21

1 Introduction

The interaction between cup products and intersection forms is a basic method to essentially analyse C^∞ -manifolds in the history (e.g., the classification theorem of simply connected manifolds). Typically, concerning the Poincaré duality with trivial coefficients, non-degeneracy of the intersection form was comprehensive in the view of the cup product. Here, it is worth noting that the interactions are reflected on algebraic futures of the coefficients $\mathbb{Z}$ and $\mathbb{Z}/p$.

However, once the (co)homology groups are investigated with local coefficients A , such an interaction has many unknown aspects with ambiguity and differences; Actually, graded commutativity and the Krull dimension of A appear as obstructions: For example, we see such a difference even from infinite cyclic cover $\tilde{Y}$ of closed manifolds; While the Blanchfield duality [Bla] on the homology of $\tilde{Y}$ is defined from some Bockstein operator to realize Hermitian intersection forms over $\mathbb{Z}[t^{\pm 1}]$, the Milnor duality [M1] is with regard to the cup products of $\tilde{Y}$ over fields and requires some assumptions; Moreover, there has been no description to explicitly connect the two duality (cf. Theorem 2.1). Furthermore, the higher duality, as in the Casson-Gordon invariants [CG] and the higher Blanchfield duality [COT] on 3-manifolds, is often approached from intersection forms of coverings; this homological approach is useful to see duality, but is not compatible with computing the duality, and with other cohomology theory as in characteristics. Further, the higher such dualities are up, the more difficulty to choose an appreciate coefficient ring we readily face, as in [COT, T1, Ka].

In this paper, we interpret three classical 3-manifold invariants from twisted pairings, and provide a method of diagrammatically computing them. The twisted product is constructed from a homomorphism $f : \pi_1(Y) \rightarrow G$, where Y is a 3-manifold, as follows: Set up a relative homology 2-class $\mu \in H_2(Y, \partial Y; \mathbb{Z})$, a right G -module M and a G -invariant bilinear function $\psi : M^{\otimes 2} \rightarrow A$ for some ring A . Then, we can define the twisted pairings as a bilinear form

$$\mathcal{Q}_\psi : H^1(Y, \partial Y; M)^{\otimes 2} \xrightarrow{\smile} H^2(Y, \partial Y; M^{\otimes 2}) \xrightarrow{\langle \bullet, \mu \rangle} M^{\otimes 2} \xrightarrow{\langle \psi, \bullet \rangle} A. \quad (1)$$

Here we regard M as the local coefficient module of Y via f , and the first map $\smile$ is the cup product, and the second (resp. third) is defined by the pairing with μ (resp. ψ). Though this pairing $\mathcal{Q}_\psi$ (1) seems speculative and uncomputable from definitions, the author [No2] has developed a diagrammatic computation of the $\mathcal{Q}_\psi$ in the link case $Y = S^3 \setminus L$ (see also §4.1).

From now on, we will explain the three 3-manifold invariants in details, where an appreciate choices of G , M and ψ is a key of the interpretation.

(I) The first one is a reduction as for the classical Blanchfield parings of knots. To be more precise (Theorem 2.1), if G is the infinite cyclic group $\mathbb{Z}$ and Y is the knot complement, the bilinear form (1) associated with appropriate choices of ψ and M is equal to the $(t + 1)/(t - 1)$ -multiple of the classical pairing; Consequently, the theorem enables us to compute the Blanchfield pairing using no Seifert surface (cf. the formulas [T2] from Seifert surfaces); As a result, we succeed in first determining the pairing of the torus knot; see Proposition 4.2. Furthermore, as a corollary, “the cocycle knot-invariants [CJKLS] from Alexander quandles” turn to be topologically characterized as the Blanchfield pairing (see Theorem B.1 for the details);

(II) Furthermore, we also address the twisted pairings of infinite cyclic covers $\tilde{Y}$, where the local coefficients arise from linear representations over a field. As in Milnor [M1, KL], the study is an approach to geometrically understand the Bockstein operator mentioned above and the (twisted) signature of knots. However, as a modification of [Neu, §11], we show (Theorem 2.2) that the cup product is reduced to the bilinear form (1) on the base space Y ; This theorem has several corollaries: First, since this description coincides with a 2-form on some twisted Alexander modules [No2, §4], we provide the 2-form a topological meaning and show a twisted version of Milnor duality [M1] (see Theorem 2.4). Furthermore, the setting as in [Go] gives an interpretation from Goldman's symplectic structure on a flat moduli space of Y (see Appendix). Moreover, since the theorem gives a diagrammatic computation of the twisted pairings in link cases, we give examples of some computations (§4.1).

(III) The third one is concerning the Casson-Gordon local signature [CG], $\text{Sign}_\omega(K, \chi)$; The signature depends on $\omega \in \mathbb{R}/\mathbb{Z}$, and is known to be a useful method to estimate the slice 4-genus of knots, as a metabelian viewpoint; see [COT]. However, to compute the signature from definitions, we readily encounter difficulties to discuss the intersection form of an appreciate 4-manifold and the twist by ω ; the known computations [CG, Gi] are only those with $\omega = 0$ of some knots. Following the idea in [KL, §6], this paper succeeds in giving a diagrammatic algorithm to practically compute the difference $\text{Sign}_\omega(K, \chi) - \text{Sign}_1(K, \chi)$; see §4.4 for the detail. Since the point is a reduction [Neu] of the intersection form to some cup products of 3-manifolds, we need describe no 4-manifold in the algorithm. However, in this paper, we do not find new examples to detect slice 4-genus; it is a problem for the further.

The three applications justify some concrete importance of the twisted pairings (1) as a setting; In summary, it is sensible to hope that the bilinear form can recover other knot invariants and more computations with appreciate choices of G , M and ψ .

This paper is organized as follows. In application, Section 2 discusses a relation to classical knot invariant, including the Blanchfield pairing of a knot and the twisted cup products of infinite cyclic coverings. Section 3 studies the the Casson-Gordon local signature. Section 4 reviews the computation in [No2], and gives some examples of the computation. Section 5 completes the proofs of the theorems.

Notation. Every knot K is a smooth, oriented and embedded of a circle in the 3-sphere S^3 . We often regard the complementary $S^3 \setminus K$ as the 3-manifold which is obtained from S^3 by removing an open tubular neighborhood of K .

2 Result I: two relations to infinite cyclic coverings

From the viewpoint of the bilinear twisted pairings $\mathcal{Q}_\psi$ in (1), we will observe some relations to classical knot invariants, including the classical Blanchfield pairing and the twisted Milnor signatures. The proofs of all the statements in Sections 2–3 will appear in §5.

2.1 Abelian case; recover of the Blanchfield pairing of knots

As the simplest case, we begin by dealing with the bilinear forms $\mathcal{Q}_\psi$ when $G = \mathbb{Z}$.

We roughly review the Blanchfield pairing [Bla] of a knot K with Alexander polynomial Δ_K . Let f be the abelianization $\pi_1(S^3 \setminus K) \rightarrow \mathbb{Z}$, and choose a generator t of $\mathbb{Z}$. The first homology $H_1(S^3 \setminus K; \mathbb{Z}[t^{\pm 1}])$ with local coefficients is called *the Alexander module*, i.e., the first homology of the infinite cyclic covering space $\tilde{Y}$ associated with f . Then, from the view of intersection forms in $\tilde{Y}$, the Blanchfield pairing is defined as a sesquilinear form

$$\text{Bl}_K : H_1(S^3 \setminus K; \mathbb{Z}[t^{\pm 1}])^{\otimes 2} \longrightarrow \mathbb{Z}[t^{\pm 1}]/(\Delta_K), \quad (2)$$

(§5.2 reviews the definition¹). This Bl_K is known to be non-singular, hermitian and isometry (see [Bla, Ka, Hil]); Further, it is a complete invariant of “the S -equivalence equivalences” in knots (see [T2, NS] for details).

Then, the following theorem asserts that the Blanchfield pairing up to constant multiples can be recovered from some bilinear form $\mathcal{Q}_\psi$:

Theorem 2.1. *Let D be a diagram of a knot K with $\#K = 1$ and Alexander polynomial Δ_K , and let M be the quotient module $\mathbb{Z}[t^{\pm 1}]/(\Delta_K)$. Define $\psi_0 : M \otimes M \rightarrow \mathbb{Z}[t^{\pm 1}]/(\Delta_K)$ by $\psi_0(x, y) = \bar{x}y$. Then, there is a $\mathbb{Z}[t^{\pm 1}]$ -module isomorphism*

$$H_1(S^3 \setminus K; \mathbb{Z}[t^{\pm 1}]) \cong H^1(S^3 \setminus K, \partial(S^3 \setminus K); M)$$

such that, for any $x, y \in H_1(S^3 \setminus K; \mathbb{Z}[t^{\pm 1}])$, the following equality holds as bilinear forms:

$$\mathcal{Q}_{\psi_0}(x, y) = \frac{1+t}{1-t} \cdot \text{Bl}_K(x, y) \in \mathbb{Z}[t^{\pm 1}]/(\Delta_K),$$

We note that $1-t$ is invertible in $\mathbb{Z}[t^{\pm 1}]/(\Delta_K)$ because of $\Delta_K(1) = \pm 1$.

In summary, we now emphasize novel advantages of this theorem. First, noting from Remark 2.3 that $\mathcal{Q}_{\psi_0}$ is equivalent to the cup product $\smile_{KL}$ on $\tilde{Y}$ considered by Milnor [M1], Theorem 2.1 consequently implies a close equality between this $\smile_{KL}$ and the pairing Bl_K in the knot case. Here the constant $(1+t)(1-t)^{-1}$ is the key to connect the hermitian pairing Bl_K with the anti-one $\mathcal{Q}_{\psi_0}$. Furthermore, the pairing Bl_K turns out to be diagrammatically computable via $\mathcal{Q}_{\psi_0}(x, y)$; see, e.g., §4.2 for a short computation on the torus knots.

2.2 Recover of twisted cup products of infinite cyclic covers

Furthermore, we will see that the pairing $\mathcal{Q}_\psi$ includes the twisted cup products (of infinite cyclic covers) considered by Milnor [M1, §4] and Kirk-Livingston [KL, §7], and show a non-degeneracy of the bilinear map. More generally, this subsection deals with not only links in S^3 but also connected compact 3-manifolds Y with boundaries ∂Y (possibly $\partial Y = \emptyset$).

To this end, we begin by briefly reviewing the observation in [KL, §7]. Take two homomorphisms $\alpha : \pi_1(Y) \rightarrow F = \mathbb{Z}$ and $f_{\text{pre}} : \pi_1(Y) \rightarrow GL_n(\mathbb{F})$ with some commutative field $\mathbb{F}$. Let $\tau : \tilde{Y} \rightarrow \tilde{Y}$ be the covering transformation. Denote the local system $\mathbb{F}^n$ by V , and take

¹In many cases (see [COT, Ka, T2, Hil]), the image is described as the universal injective module $\mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]$. However, such as [T1], the pairing factors through the inclusion $\mathbb{Z}[t^{\pm 1}]/(\Delta_K) \hookrightarrow \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]$ sending $[f]$ to $[f/\Delta_K]$.

a $\pi_1(Y)$ -invariant bilinear function $\psi_{\text{pre}} : V \otimes V \rightarrow \mathbb{F}$. Let $\mu_\alpha \in H_2(\tilde{Y}, \partial\tilde{Y}; \mathbb{Z})$ be a lift of the dual element of $[\alpha] \in H^1(Y; \mathbb{Z})$. In this situation, Kirk and Livingston considered the bilinear map $\smile_{KL}$ defined by setting the composite

$$\smile_{KL} : H^1(\tilde{Y}, \partial\tilde{Y}; V)^{\otimes 2} \xrightarrow{\sim} H^2(\tilde{Y}, \partial\tilde{Y}; V \otimes V) \xrightarrow{\langle \psi_{\text{pre}}, \bullet \rangle} H^2(\tilde{Y}, \partial\tilde{Y}; \mathbb{F}) \xrightarrow{\cap \mu_\alpha} \mathbb{F}. \quad (3)$$

Further, they claimed without proof a non-degeneracy [KL, Theorem 7.1] under a certain assumption with $\mathbb{F} = \mathbb{C}$. Therefore, if $\mathbb{F}$ is the complex field $\mathbb{C}$, for $z \in \mathbb{Z}$, we can define the signature of $\smile_{KL}$ on the $(\tau^* - z)$ -primary invariant part of $H^1(\tilde{Y}, \partial\tilde{Y}; V)$. This signature valued in $\mathbb{Z}$ is called the *twisted Milnor signature*; see [M1, §5] for the original one. Further, the authors indicated a connection between this twisted signature and the Casson-Gordon signature, although they gave no examples and no computation, and did not explicitly describe the relation in details. Actually, this $\smile_{KL}$ also seems uncomputable by definition.

We will reformulate (Theorem 2.2) the above bilinear form $\smile_{KL}$ in terms of the bilinear form (1). For this, define a right $\mathbb{F}[\pi_1(Y)]$ -module $V[t^{\pm 1}] := \mathbb{F}[t^{\pm 1}] \otimes_{\mathbb{F}} V$ by taking the tensor product $\bar{\alpha} \otimes f_{\text{pre}}$; the action is given by

$$(q \otimes v) \cdot g := (qt^{\alpha(g)}) \otimes (v \cdot f_{\text{pre}}(g)), \quad \text{for } g \in \pi_1(Y), \quad q \in \mathbb{F}[t^{\pm 1}], \quad v \in V.$$

Further, we suppose a common assumption similar to [KL, HSW, Hil].

Assumption ($\dagger$). Suppose that R is a field $\mathbb{F}$ of characteristic zero with involution $-$. Furthermore, there exist $y_0 \in \pi_1(Y)$ and $d_0 \in \pi_1(\partial Y)$ with $\alpha(y_0) = \alpha(d_0) = 1$ as above. Moreover, the twisted first homology $H_1(Y, \partial Y; V[t^{\pm 1}]^n)$ is of finite dimension over $\mathbb{F}$, that is, its order or the twisted Alexander polynomial, Δ_f , is non-zero: $\Delta_f \neq 0$. Further, Δ_f is invertible in the zeroth homology $H_0(Y, \partial Y; V[t^{\pm 1}]^n)$.

Furthermore, we introduce a trace map similar to [T2, §2] and [Neu, §11]. According to $\pm$, take the two rings of formal Laurent series, which are defined as

$$\mathbb{F}[[\mathbb{Z}]]_{\pm} := \{ \sum a_i t^i \in \mathbb{F}[[t]] \mid a_{\pm k} = 0 \text{ for } k \text{ sufficiently small.} \}.$$

Since every $x \neq 0$ in $\mathbb{F}[t^{\pm 1}]$ is invertible in $\mathbb{F}[[\mathbb{Z}]]_{\pm}$, we have a natural injection $i_{\pm} : \mathbb{F}(t) \hookrightarrow \mathbb{F}[[\mathbb{Z}]]_{\pm}$. For $x \in \mathbb{F}(t)$, define $\text{tr}(x)$ to be $i_+(x)|_{t^0} - i_-(x)|_{t^0}$, where the subscript $|_{t^0}$ means the coefficient of t^0 . Then, this map induces a well-defined $\mathbb{F}$ -linear map

$$\text{Tr} : \mathbb{F}[t^{\pm 1}]/(\Delta_f) \hookrightarrow \mathbb{F}(t)/\mathbb{F}[t^{\pm 1}] \xrightarrow{\text{tr}} \mathbb{F}; \quad x \mapsto [x/\Delta_f] \mapsto \text{tr}([x/\Delta_f]).$$

We can easily verify that, if Δ_f is irreducible, Tr equals the trace map from the field extension.

Theorem 2.2. *Let the coefficients $V = \mathbb{F}^n$ and $M = V \otimes_{\mathbb{F}} \mathbb{F}[t^{\pm 1}]/(\Delta_f)$ satisfy in Assumption ($\dagger$). Define $\psi : M^2 \rightarrow \mathbb{F}[t^{\pm 1}]/(\Delta_f)$ by $\psi(at^i, bt^j) := t^{i-j} \psi_{\text{pre}}(a, b)$ for $a, b \in V$.*

Then, there is a canonical $\mathbb{F}$ -isomorphism $\Upsilon_{\Delta}^* : H^1(\tilde{Y}, \partial\tilde{Y}; V) \cong H^1(Y, \partial Y; M)$, for which the equality $\smile_{KL} \circ (\Upsilon_{\Delta}^*)^{\otimes 2} = \text{Tr} \circ \mathcal{Q}_{\psi} : H^1(Y, \partial Y; M)^{\otimes 2} \rightarrow \mathbb{F}$ holds: Precisely,

$$\begin{array}{ccc} H^1(Y, \partial Y; M)^{\otimes 2} & \xrightarrow{\mathcal{Q}_{\psi}} & \mathbb{F}[t^{\pm 1}]/\Delta_f \\ \downarrow (\Upsilon_{\Delta}^*)^{\otimes 2} \cong & \circlearrowleft & \downarrow \text{Tr} \\ H^1(\tilde{Y}, \partial\tilde{Y}; V)^{\otimes 2} & \xrightarrow{\smile_{KL}} & \mathbb{F}. \end{array} \quad (4)$$

Remark 2.3. Conversely, similar to [Neu, Remark in §11], the previous bilinear form $\mathcal{Q}_\psi$ recovers from $\smile_{KL}$. To be precise, we can easily see $[a] = [i_+^{-1}(\sum_{j \geq 0} \text{tr}(t^{-j}a) \cdot t^j)]$ for all $[a] \in \mathbb{F}(t)/\mathbb{F}[t^{\pm 1}]$ (cf. Laurent expansion); therefore, $[\mathcal{Q}_\psi(x, y)/\Delta_f] = [i_+^{-1}(\sum_{j \geq 0} \Upsilon_\Delta^*(\tau_*^{-j}x) \smile_{KL} \Upsilon_\Delta^*(y) \cdot t^j)]$. In summary, our formula $\mathcal{Q}_\psi$ and the bilinear form $\smile_{KL}$ are equivalent.

As a result, if Y is a link component $S^3 \setminus L$, Theorem 4.1 ensures a diagrammatic computation of $\smile_{KL}$; see §4.3 for examples.

Furthermore, let us mention a non-degeneracy of $\mathcal{Q}_\psi$ (Theorem 2.4). The *dual* (or *conjugredient*) representation, f_{pre}^* , of f_{pre} is defined by $f_{\text{pre}}^*(g)^{-1} = \overline{f_{\text{pre}}(g)}$. It is known (see [HSW, Lemma 3.1]) that there exists a bilinear form $\psi_{\text{pre}} : \mathbb{F}^n \otimes \mathbb{F}^n \rightarrow \mathbb{F}$ which is non-degenerate if and only if the original representation f_{pre} is conjugate to the dual one f_{pre}^* . Thus, the symplectic or unitary representations of $\pi_1(Y)$ are excellent examples. In addition, consider the connecting map $\delta^* : H^0(\partial Y; M) \rightarrow H^1(Y, \partial Y; M)$, and denote the cokernel by $\overline{H^1(Y, \partial Y; M)}$. Since the restriction of $\mathcal{Q}_\psi$ on $\text{Im}(\delta^*)$ is zero, the non-degeneracy is stated as follows:

Theorem 2.4. *Put the previous quotient $\overline{H^1(Y, \partial Y; M)}$. As in notation in Theorem 2.2, if there is a bilinear form $\psi_{\text{pre}} : \mathbb{F}^n \otimes \mathbb{F}^n \rightarrow \mathbb{F}$ which is $\pi_1(Y)$ -invariant and is non-degenerate, then the associated bilinear form $\mathcal{Q}_\psi$ (4) on the quotient $\overline{H^1(Y, \partial Y; M)}^{\otimes 2}$ is non-degenerate.*

Remark 2.5. As in [M1], the assumption $\text{Char } \mathbb{F} = 0$ is important. For example, Theorem 2.1 implies that, even if the coefficient $\mathbb{F}$ is trivial and Y is a knot complement, the duality does not hold on the non-zero integer $\Delta(-1) \in \mathbb{Z}$.

As is classically known, the Blanchfield duality implies reciprocity of Alexander polynomials. In analogy, we immediately obtain from Theorem 2.4 a known criterion for reciprocity of the twisted polynomial Δ_f .

Corollary 2.6 (cf. [HSW, Theorem 3.2]). *Let the coefficients $V = \mathbb{F}^n$ and $M = V \otimes_{\mathbb{F}} \mathbb{F}[t^{\pm 1}]/(\Delta_f)$ be as in Assumption (†) and Theorem 2.4. Then the polynomial Δ_f has reciprocity. Precisely, there are $N \in \mathbb{Z}$ and $a \in \mathbb{F}^\times$ so that*

$$\Delta_f = at^N \overline{\Delta_f}.$$

Here, we should refer the reader to [HSW, FKK], which say the same statement in the knot case. Their proof was an approach from a Reidemeister torsion, and later developed as a general reciprocity; see [FKK].

To sum up, we mention the known fact (see [M2, §3]) that, if ψ_{pre} is (skew)hermitian and the base space have reciprocity, the isometry classes of such bilinear forms have been completely classified. Here, the classification is done by using the trace maps in extension fields. So, according to the complete invariants stated in [M2, §3], we can deal with concretely the bilinear form $\mathcal{Q}_\psi$.

3 Result II; Recover of the Casson-Gordon local signature

The purpose of this section is to state the recover. However, in contract to the previous section, we have to prepare several terminologies in §3.1.

3.1 Review; the Casson-Gordon invariants and σ -signature

We begin by recalling contexts for the Casson-Gordon invariants. Let K be a tame knot embedded in the 3-sphere S^3 with orientation. For prime powers q and $d \in \mathbb{Z}$, we denote by B_q the q -fold cyclic cover branched along K , and write E_q for the q -fold cyclic cover of $S^3 \setminus K$. Let X_q denote the q -fold cyclic cover of 0-framed surgery on K . Then the inclusions $i : E_q \hookrightarrow X_q$ and $j : E_q \hookrightarrow B_q$ induce an isomorphism $i_* : H_1(E_q; \mathbb{Z}) \cong H_1(X_q; \mathbb{Z})$ and a surjection $j_* : H_1(E_q; \mathbb{Z}) \rightarrow H_1(B_q; \mathbb{Z})$, respectively. Thus, we have a surjection

$$j_* \circ (i_*)^{-1} : H_1(X_q; \mathbb{Z}) \longrightarrow H_1(B_q; \mathbb{Z}).$$

Since $\mathbb{Z}/d$ acts on $\mathbb{Q}(\zeta_d)$ with $\zeta_d = \exp(2\pi\sqrt{-1}/d)$ canonically, we fix an inclusion $\mathbb{Z}/d \hookrightarrow GL_1(\mathbb{C})$ in what follows. Then, a given character $\chi : \pi_1(B_q) \rightarrow \mathbb{Z}/d$ defines a representation

$$\bar{\chi} : \pi_1(X_q) \longrightarrow GL_1(\mathbb{Q}(\zeta_d)) \hookrightarrow GL_1(\mathbb{C})$$

as the composite of the surjection $j_* \circ (i_*)^{-1}$ and the Hurewicz map of X_q . Further, the abelianization $\alpha : \pi_1(S^3 \setminus K) \rightarrow \mathbb{Z}$ yields $\bar{\alpha} : \pi_1(X_q) \rightarrow \mathbb{Z} = \langle t \rangle$. Then, by the basic fact in bordism theory (see, e.g., [FG, §2.2]), there is an oriented compact 4-manifold W such that the boundary is the q -copies of X_q and that $\bar{\chi} \times \bar{\alpha}$ extends componentwise to $\hat{\chi}_W \times \hat{\alpha} : H_1(W; \mathbb{Z}) \rightarrow \mathbb{Z}/d \times \mathbb{Z}$ via the boundary injection. To summarize, here is a description of the manifolds appearing above:

$$\begin{array}{ccccc} B_q & \xleftarrow{\quad} & E_q & \xrightarrow{\text{0-surgery}} & X_q & X_q \# \cdots \# X_q = \partial W \subset W \\ \text{Branched covering} \downarrow & & \downarrow q\text{-fold covering} & & \downarrow \text{Branched covering} \\ S^3 & \xleftarrow{\quad} & S^3 \setminus K & \xrightarrow{\text{0-surgery}} & X_K \end{array}$$

Furthermore, take the trivial homomorphism $\chi_{\text{tri}} : \pi_1(W) \rightarrow GL_1(\mathbb{Q})$. Then, we have the three cohomologies of W with local coefficient rings $\mathbb{Q}$, $\mathbb{C}$ and $\mathbb{C}(t)$ via χ_{tri} , $\hat{\chi}_W$ and $\hat{\chi}_W \otimes \bar{\alpha}$, respectively. Since $H_2(W; \mathbb{C}(t))$ is known to be of finite dimension over $\mathbb{C}(t)$; see [CG, Gi], we accordingly obtain the respect intersection forms I_{tri} , $I_{\hat{\chi}}$ and $I_{\bar{\alpha} \otimes \hat{\chi}}$ on the space $H_2(W; \dagger)$, which are known to be non-degenerate hermitian (see, e.g., [FG, §2] for the detailed formula).

Then, the *Casson-Gordon invariant* is defined as the following element:

$$\text{CG}(K, \chi) := (I_{\bar{\alpha} \otimes \hat{\chi}} - I_{\text{tri}}) \otimes (1/q) \in \text{Witt}(\mathbb{C}(t)) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q],$$

where $\text{Witt}(\mathbb{C}(t))$ is the (hermitian) Witt group of the functional field $\mathbb{C}(t)$. We can easily see their independence from the choices of W and $\hat{\chi}_W$, as in [CG, Neu, COT]. Further, roughly speaking, if K and χ satisfy certain conditions, $\text{CG}(K, \chi)$ is an additively concordance invariant (see [CG, Gi, COT] for the details).

Further, as a reduction of the origin $\text{CG}(K, \chi)$, it is an easy way to consider the discriminant (see [KL, HKL]) or the (hermitian) local signature as reductions of $\text{CG}(K, \chi)$. This paper examines the latter. To this end, let us call a complex number $w \in \mathbb{C}$ *generic*, if w is not a solution of $\det(I_{\bar{\alpha} \otimes \hat{\chi}}) = 0 \in \mathbb{C}(t)$ and $t - w$ is invertible in $H_2(W; \mathbb{C}(t))$. By definition, non-generic points are finitely many in $\mathbb{C}$. Moreover, if $w \in \mathbb{C}$ is generic and $|w| = 1$, we can define

the signature $\text{Sign}_w(\text{CG}(K, \chi)) \in \mathbb{Z}[1/q]$; By definition, $\text{Sign}_w(\text{CG}(K, \chi))$ is locally constant on the circle $S^1 \subset \mathbb{C}$ by removing the non-generic points. However, the signatures at non-generic points have some difficulties. For example, concerning $w = 1$, consider $\sigma_1(K, \chi) \in \mathbb{Q}$ defined to be $(\text{Sign}(I_{\hat{\chi}}) - \text{Sign}(I_{\text{tri}}))/q$, as in [CG]. Then $\sigma_1(K, \chi) - \lim_{t \rightarrow +0} \text{Sign}_{\exp(\sqrt{-1}t)}(\text{CG}(K, \chi))$ is called *nullity*, and is not always zero together with a relation to the slice-ribbon conjecture (see [CG, FG]). See the computation with $w = 1$ in [CG, KL, FG] and therein references

However, the definition of $\text{CG}(K, \chi)$ in the first place requires such a 4-manifold W and an extension $\hat{\chi}_W$; in the very beginning, it has been considered to be hard to compute the local signatures.

3.2 A diagrammatic formulation

We will state the main theorem 3.1 for computing the related signature from a knot diagram and cup products. We denote $\pi_1(S^3 \setminus K)$ by π_K , for short.

We begin by setting up a map f_χ and a module M as follows. Since the abelianization $\pi_K \rightarrow \mathbb{Z}$ splits, we choose an isomorphism $\pi_K \cong \pi'_K \rtimes \mathbb{Z}$. Since $\pi_1(E_q)$ injects the commutator subgroup π'_K , the composite $\hat{\chi} \circ i_*^{-1} : \pi_1(E_q) \rightarrow GL_1(\mathbb{C})$ gives rise to the induced representation $f_\chi : \pi_K \rightarrow GL_q(\mathbb{C}[t^{\pm 1}])$. To be precise, f_χ is easily remembered in the matrix presentation

$$f_\chi(g, n) := \begin{pmatrix} \zeta_d^{\hat{\chi}(g)} & 0 & \cdots & 0 \\ 0 & \zeta_d^{\hat{\chi}(g \cdot 1)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \zeta_d^{\hat{\chi}(g \cdot (q-1))} \end{pmatrix} \cdot \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & \ddots & 1 \\ t & 0 & \cdots & \cdots & 0 \end{pmatrix}^n \in GL_q(\mathbb{C}[t^{\pm 1}]), \quad (5)$$

for $(g, n) \in \pi'_K \rtimes \mathbb{Z}$. Since it is known [CG, Gi] that the twisted homology $H_1^t(S^3 \setminus K; (\mathbb{C}[t^{\pm 1}])^q)$ is of finite dimension, we have the non-zero twisted Alexander polynomial $\Delta_\chi \in \mathbb{C}[t^{\pm 1}]$ with the reciprocity $\Delta_\chi = bt^n \overline{\Delta_\chi}$ for some $n \in \mathbb{Z}$, $b \in \mathbb{C}^\times$ (see Theorem 2.4). Then, we let M be the quotient module $(\mathbb{C}[t^{\pm 1}]/\Delta_\chi)^q$ with involution, and let π_K act canonically on M via f_χ .

Let us turn into reviewing briefly notions in [Neu, §2-3], in order to apply the related signature to [Neu, Theorem 3.2]. In general, given a finite dimensional $\mathbb{C}$ -vector space V acted on by $\mathbb{Z} = \langle \tau_* \rangle$, for $z \in \mathbb{C}$, let us consider the $(\tau_* - z)$ -primary invariant subspace, and denote it by V_z , i.e., $V_z := \{v \in V \mid v(\tau_* - z)^k = 0 \text{ for some } k\}$. Further, if V has a skew hermitian bilinear form b over $\mathbb{C}$, we define a value $\lambda(V, b) \in \mathbb{R}$ to be $\sum_{z: |z|=1} \lambda(V_z, b)$, and

$$\lambda(V_z, b) = \begin{cases} (2a - 1) \cdot \text{Sign}(V_z, \sqrt{-1}b), & z = \exp(2\pi\sqrt{-1}a) \text{ with } 0 < a < 1, \\ -\text{Sign}(V_1, \sqrt{-1}b_{(1)}), & z = 1, \end{cases}$$

where $b_{(1)}$ is defined by $b_{(1)}(x, y) = b(\tau_* x - \tau_*^{-1} x, y)$.

We will describe the main theorem by applying the situations in §3.2 to the previous settings. Denote the cohomology $H^1(S^3 \setminus K, \partial S^3 \setminus K; M)$ by $\mathcal{H}$. Given a character χ , combining $\mathcal{Q}_\psi$ in (1) with the trace map in §2.2 gives a skew-hermitian 2-form $b := \text{Tr} \circ \mathcal{Q}_\psi$ on $\mathcal{H}$. Notice that $\mathcal{H}$ is of finite dimension over $\mathbb{C}$, and is acted on by t . Further, for $w \in \mathbb{C}$, define the map $\hat{\alpha}_w : \pi_1(X_q) \rightarrow \mathbb{C}$ by setting $\gamma \mapsto w^{\alpha(\gamma)}$, where α is the abelianization of $\pi_1(X_q)$. Tensoring with $\mathbb{C}$ via $\hat{\alpha}_w$, we can twist the action of $\mathbb{Z}$ on $\mathcal{H} \otimes_{\mathbb{C}} \mathbb{C}$. Denote the $\mathbb{C}$ -space with action by $\mathcal{H} \otimes \tau$, for short. To summarize, here is the required statement:

Theorem 3.1. *Assume the following four conditions: (i) the homomorphism χ is non-trivial. (ii) $w \in \mathbb{C}$ is generic. (iii) Δ_χ is not divisible by $t - \omega$. (iv) Δ_χ is not divisible by $1 - t$.*

Then, the following equality holds:

$$\text{Sign}_w(K, \chi) - \sigma_1(K, \chi) = \lambda(\mathcal{H}, b) - \lambda(\mathcal{H} \otimes \tau, b \otimes w) \in \mathbb{Z}[1/q].$$

While we defer the proof until §5.3, we now sum up the theorem. We can diagrammatically compute the right hand side following from Proposition 3.1, and, consequently can do the left one. Needless to say, in order to compute the goal $\text{Sign}_w(K, \chi)$, we have to investigate the second term $\sigma_1(K, \chi)$ and a condition of χ .

Finally, we emphasize that the four conditions in Theorem 3.1 is not strong. Actually, (i) is obviously weak, and there are finitely many w in $\mathbb{C}$ that satisfy neither (ii) nor (iii). In contract, Δ_χ does not always satisfy (iv). However, as seen in the proof, the condition (iv) arises from a Mayer-Vietoris discussion: Thus, even when Δ_χ is divisible by $1 - t$, we can get a similar equality by following carefully the proof; However, we omit describing the detail, since it depends on the knot type.

4 Examples of computations

This section, by composing the theorems in the previous sections with the result in [No2]. To more precise, Section 4.1 strictly describes diagrammatic computation of the cup products, and Sections 4.2–4.4 give some examples.

4.1 Reviews; diagrammatic computation of the twisted pairings.

We will need some knowledge of quandles before proceeding.

Thorough this section, we fix a group G and a right G -module M over a ring A . Let Z be a subset of G closed under the conjugation operation. Further, denote $M \times Z$ by X , and define a binary operation on X by

$$\triangleleft : (M \times Z) \times (M \times Z) \longrightarrow M \times Z, \quad (a, g, b, h) \longmapsto ((a - b) \cdot h + b, h^{-1}gh). \quad (6)$$

Further, given a knot $K \subset S^3$ with a group homomorphism $f : \pi_1(S^3 \setminus K) \rightarrow G$ such that every meridian is sent to Z . Let us choose an oriented knot diagram D of K . Then, it follows from the conjugacy operation of G and the Wirtinger presentation of D that the homomorphism f is regarded as a map $\{\text{arcs of } D\} \rightarrow G$. Furthermore, a map $\mathcal{C} : \{\text{arcs of } D\} \rightarrow X$ over f is an X -coloring over f if it satisfies $\mathcal{C}(\alpha_\tau) \triangleleft \mathcal{C}(\beta_\tau) = \mathcal{C}(\gamma_\tau)$ at each crossings of D illustrated as Figure 1. Let $\text{Col}_X(D_f)$ denote the set of all X -colorings over f . By definition, this $\text{Col}_X(D_f)$ canonically injects into the product $M^{\alpha(D)}$, where $\alpha(D)$ is the number of arcs of D . Hence, $\text{Col}_X(D_f)$ serves as an A -submodule of $M^{\alpha(D)}$. Furthermore, the diagonal submodule $M_{\text{diag}} \subset M^{\alpha(D)}$ is contained in $\text{Col}_X(D_f)$, and is a direct summand M of $\text{Col}_X(D_f)$. Thus, we denote by $\text{Col}_X^{\text{red}}(D_f)$ another direct summand.

Furthermore, one introduces a bilinear form on the A -module $\text{Col}_X(D_f)$ as follows. Let $\psi : M^2 \rightarrow A$ be a bilinear map over $\mathbb{Z}$. Moreover, we assume that this ψ is G -invariant and is



Figure 1: Positive and negative crossings, and complementary regions marked by A .

over A . Precisely, the two equalities

$$\psi(xa, yb) = \bar{a}\psi(x, y)b, \quad \psi(x \cdot z, y \cdot z) = \psi(x, y)$$

hold for any $a, b \in A$, $x, y \in M$ and $z \in G$. We further define the two maps $\mathcal{D}_1 : X^2 \rightarrow M$ and $\mathcal{D}_2 : X \rightarrow M$ by setting $\mathcal{D}_1((x, z), (y, w)) = x - y$ and $\mathcal{D}_2(y, w) = y - y \cdot w$, respectively. Furthermore, we define a map

$$\mathcal{Q}_\psi : (\text{Col}_X(D_f))^2 \longrightarrow A; \quad (\mathcal{C}, \mathcal{C}') \longmapsto \sum_{\tau} \epsilon_{\tau} \cdot \psi(\mathcal{D}_1(\mathcal{C}(\alpha_{\tau}), \mathcal{C}(\beta_{\tau})), \mathcal{D}_2(\mathcal{C}'(\beta_{\tau}))), \quad (7)$$

where τ runs over all the crossings of D , and the symbols α_{τ} , β_{τ} are the arcs and $\epsilon_{\tau} \in \{\pm 1\}$ is the sign of τ according to Figure 1.

Next, from the view of this theorem we will reformulate the bilinear form $\mathcal{Q}_\psi$ defined in (1). To see this, we set up notation. Let M^{op} denote the A -module on M with the operation $x \cdot a := x\bar{a}$ for $x \in M, a \in A$. Letting $Y_K = S^3 \setminus K$, we should notice the isomorphism

$$H^*(Y_K, \partial Y_K; M) \cong H^*(Y_K, \partial Y_K; M^{\text{op}})$$

by definitions. In addition, recall the isomorphism $H_2(Y_K, \partial Y_K; \mathbb{Z}) \cong \mathbb{Z}$ with a basis μ which corresponds to a Seifert surface in $S^3 \setminus K$.

Theorem 4.1. *Let $Y_K = S^3 \setminus K$ and M be as above. Then, there are A -module isomorphisms*

$$\text{Col}_X(D_f) \cong H^1(Y_K, \partial Y_K; M) \oplus M, \quad \text{Col}_X^{\text{red}}(D_f) \cong H^1(Y_K, \partial Y_K; M).$$

Further, the restriction of $\mathcal{Q}_\psi$ on $\text{Col}_X^{\text{red}}(D_f)$ is equal to the following composite:

$$H^1(Y_K, \partial Y_K; M)^{\otimes 2} \xrightarrow{\sim} H^2(Y_K, \partial Y_K; M^{\text{op}} \otimes M) \xrightarrow{\langle \bullet, \mu \rangle} M^{\text{op}} \otimes M \xrightarrow{\langle \psi, \bullet \rangle} A.$$

To summarize, the point is that, given a diagram D , we can diagrammatically compute the form $\mathcal{Q}_\psi$ by definitions.

4.2 The Blanchfield pairing of the torus knot

Parallel to the setting in Theorem 2.1, we will observe the case where the subset $Z \subset G = \mathbb{Z}$ consists of the generator $\{t\}$ alone. Then the module M is a $\mathbb{Z}[t^{\pm 1}]$ -module, and the operation on $X = Z \times M = M$ in (6) is formulated by $x \triangleleft y := tx + (1 - t)y$ for $x, y \in M$. In other ward, X with the operation is exactly the so-called *Alexander quandle* (see [CJKLS, Joy]).

Thanks to Theorem 2.1, we will address the first to determine the Blanchfield pairing of the (m, n) -torus knot $T_{m, n}$. Here note the known fact that the Alexander module is isomorphic to $\mathbb{Z}[t^{\pm 1}]/(\Delta_K)$, where Δ_K is the Alexander polynomial $(t^{nm} - 1)(t - 1)/((t^n - 1)(t^m - 1))$.



Figure 2: The eight knot and the $T_{m,n}$ -torus link with labeled arcs.

Proposition 4.2. Fix $(n, m, a, b) \in \mathbb{Z}^4$ with $an + bm = 1$, and let $K = T_{m,n}$. Then,

$$\text{Bl}_K(y_1, y_2) = \frac{nm}{(1+t^{-1})(1-t^{bm})(1-t^{an})} \cdot \bar{y}_1 y_2 \in \mathbb{Z}[t^{\pm 1}]/(\Delta_K),$$

for $y_1, y_2 \in H_1(S^3 \setminus T_{m,n}; \mathbb{Z}[t^{\pm 1}]) \cong \mathbb{Z}[t^{\pm 1}]/(\Delta_K)$.

Remark that the coefficient in this formula lies in $\mathbb{Z}[t^{\pm 1}]/(\Delta_K)$ because of l'Hôpital's rule in $\mathbb{Z}[t^{\pm 1}]$. Furthermore, it seems hard to compute the pairing from the previous formula [T2, Proposition 2.5] using the Seifert matrix, since the Seifert genus is $(n-1)(m-1)/2$; for example, Matsumoto [Mat] uses a technical formula to compute all the local signatures of $T_{m,n}$. However, our result is still simpler as follows:

Proof. Let $X = A = \mathbb{Z}[t^{\pm 1}]/(\Delta_K)$. By Theorem 2.1, we have $\text{Col}_X(D) \cong H_1(S^3 \setminus K; A) \oplus X \cong X^2$; It is enough to compute the bilinear form $\mathcal{Q}_\psi$ with $\psi(y, z) = \bar{y}z$.

To this end, let us start by examining $\text{Col}_X(D)$ in more details. Let $\alpha_1, \dots, \alpha_m$ be the arcs depicted in Figure 2. Because of the shape of D , every coloring in $\text{Col}_X(D)$ is characterized by colors of these n arcs. Hence, we can view $\text{Col}_X(D)$ as a submodule of X^m . In addition, consider elements of the forms

$$\vec{y}_k = (\delta_k, y_k + \delta_k, \frac{1-t^{2an}}{1-t^{an}}y_k + \delta_k, \dots, \frac{1-t^{an(m-1)}}{1-t^{an}}y_k + \delta_k) \in X^m, \quad (8)$$

for some $y_k, \delta_k \in X$. From this view, we can easily see that these elements (8) define X -colorings. Further, by observing the first and second components, these elements give a basis of $\text{Col}_X(D) \cong X^2$. Let $\tau_{i,j}$ be the j -th crossing point on the arc α_i . Then, concerning the X -coloring $\mathcal{C}$ arising from (8), the colors around $\tau_{i,j}$ as Figure 1 are formulated as

$$(\mathcal{C}(\alpha_{\tau_{i,j}}), \mathcal{C}(\beta_{\tau_{i,j}})) = (\frac{1-t^{bmj}}{1-t^{bm}}t^{an(i-2)}y_k + \frac{1-t^{an(i-1)}}{1-t^{an}}y_k + \delta_k, \frac{1-t^{an(i-1)}}{1-t^{an}}y_k + \delta_k) \in A^2.$$

Accordingly, we now deal with the 2-form $\mathcal{Q}_\psi(\vec{y}_1, \vec{y}_2)$. By definition, compute it as

$$\begin{aligned} & \sum_{i \leq m, j \leq n-1} \psi \left(t^{an(i-2)} \frac{1-t^{bmj}}{1-t^{bm}} y_1, \left(\frac{1-t^{an(i-1)}}{1-t^{an}} y_2 + \delta_2 \right) (1-t^{-1}) \right) \\ &= \psi \left(\sum_{j=1}^{n-1} \frac{1-t^{bmj}}{1-t^{bm}} y_1, \left(\sum_{i=1}^m \frac{t^{an} - t^{an(2-i)}}{1-t^{an}} y_2 (1-t^{-1}) \right) \right) \\ &= \psi \left(\frac{1-t^{bm(n-1)} - (n-1)(1-t^{bm})}{(1-t^{bm})^2} y_1, \frac{m(1-t^{an}) + 1-t^{-am n}}{(1-t^{an})^2} t^{an} \cdot y_2 (1-t^{-1}) \right) \end{aligned}$$

$$= \psi\left(\frac{n}{1-t^{bm}} \cdot y_1, t^{an} \frac{m}{1-t^{an}} \cdot y_2(1-t^{-1})\right) = \frac{nm(1-t^{-1})}{(1-t^{bm})(1-t^{an})} \cdot \bar{y}_1 y_2 \in \mathbb{Z}[t^{\pm 1}]/(\Delta_K),$$

where the first equality is obtained by the t -invariance of ψ and the equality $\sum_{i=1}^m t^{ani} = 0$, and an elementary computation lifted in $\mathbb{Z}[t^{\pm 1}]$ can imply the third equality by noting $t^{mn} = 1 \in X$. The last term is $(1-t)^{-1}(1+t)\text{Bl}_K(y_1, y_2)$ by Theorem 2.1. So, notice that, $\Delta_K(-1) = 1$ if nm is odd, and $\Delta_K(-1) = m$ if n is even; we may divide the value $\mathcal{Q}_\psi(\vec{y}_1, \vec{y}_2)$ by $1+t$. In the sequel, since $(1-t)^{-1} \in A$, the form $\mathcal{Q}_\psi(\vec{y}_1, \vec{y}_2)$ is reduced to the required formula. $\square$

4.3 The figure eight knot

We will compute the cup product $\smile_{KL}$ in (3) of the figure eight knot as illustrated in Figure 2. Here, we set $G = SL_2(\mathbb{F})$, where $\mathbb{F}$ is a field of characteristic 0.

Let set up the situation. Let $a_i \in \pi_1(S^3 \setminus K)$ be the element corresponding with the arc α_i in Figure 2. Then, we will employ the elliptic representation $f : \pi_1(S^3 \setminus K) \rightarrow SL_2(\mathbb{F}[t^{\pm 1}])$ such that

$$f(a_1) = t \cdot \begin{pmatrix} s & 1 \\ 0 & s^{-1} \end{pmatrix}, \quad f(a_2) = t \cdot \begin{pmatrix} s & 0 \\ u+1 & s^{-1} \end{pmatrix},$$

for some $s, u \in \mathbb{F}^\times$ with $\bar{s} = s$ and $\bar{u} = u$. It is known that s and u must satisfy $s^2 + s^{-2} + u + u^{-1} - 1 = 0$, and that the set $\text{Hom}(\pi_1(S^3 \setminus K), G)$ is characterized by such f 's.

Let us review the computation [No2, §3.2] of $\mathcal{Q}_\psi$, without the proofs. First, for simplicity and the assumption $(\dagger)$, let us suppose a generic assumption of the form

$$(u-1)(u+u^{-1}-1)(2u+2u^{-1}-1)(2u+2u^{-1}-5)(u^3-u^2-2u-1) \neq 0.$$

In addition, inspired by [Go], we discuss $\mathcal{Q}_\psi$ associated with adjoint representations; Consider the lie algebra $\mathfrak{g} = \{B \in \text{Mat}(2 \times 2; \mathbb{F}) \mid \text{Tr} B = 0\}$ with adjoint action of $SL_2(\mathbb{F})$. Then, the twisted Alexander polynomial Δ_f is equal to $t^2 + (2u + 2u^{-1} - 3)t + 1$. Thus, setting M by $\mathfrak{g}[t^{\pm 1}]/(\Delta_f)$, the cohomology $H^1(Y, \partial Y; M)$ is shown to be $\mathbb{F}[t]/(\Delta_f)$. Furthermore, taking the Killing 2-form $\psi : \mathfrak{g}^2 \rightarrow \mathbb{F}$ that takes (X, Y) to $\text{Tr}(XY)$, one showed that

$$\mathcal{Q}_\psi(\vec{x}, \vec{x}') = 2(t-t^{-1})(u+u^{-1}-1)(1-u)(u^3-u^2-2u-1)x\bar{x}'. \quad (9)$$

We will compute the cup product $\smile_{KL}$ in (3). It can be easily seen that $\text{Tr}(t-t^{-1}) = 2(u+u^{-1}-3)$ by definition. Then, Theorem 2.2 implies the conclusion

$$x \smile_{KL} y = \text{Tr} \circ \mathcal{Q}_\psi(x, y) = 4(u+u^{-1}-3)(u+u^{-1}-1)(1-u)(u^3-u^2-2u-1)xy,$$

where x, y are contained in $H^1(\tilde{Y}, \partial Y; \mathbb{F}^2) \cong \mathbb{F}[t]/(\Delta_f)$.

4.4 A computation of the Casson-Gordon local signature

In this section, we give a computation of the local signature of the knot 12_{a169} ; see Figure 3. Although the knot is known to be non-slice, we now prove the fact by using the local signature.

The computation on the twisted Alexander polynomial is based on [HKL, §10.2]. Let us review the computation. In what follows, we fix $q = 3$, and $d = 5$, and denote by ζ




Figure 3: The knot 11_{a169} with labeled arcs.

denote the fifth root of unity $\exp(2\pi\sqrt{-1}/5)$. Then, $H_1(B_q; \mathbb{Z})$ is known to be $\mathbb{Z}/25 \oplus \mathbb{Z}/25$; Further, $H_1(B_q; \mathbb{Z}/5)$ is $\mathbb{F}_5[t^{\pm 1}]$ -isomorphic to $\mathbb{F}_5[t]/(1+t+t^2)$. Therefore, for every map $\chi : \pi_1(B_q) \rightarrow \mathbb{Z}/5$ which is not trivial, the associated twisted Alexander polynomial is the same. So we fix such a χ , and the associated homomorphism defined from

$$f_\chi(\alpha_1) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ t & 0 & 0 \end{pmatrix}, \quad f_\chi(\alpha_2) = \begin{pmatrix} 0 & \zeta^2 & 0 \\ 0 & 0 & \zeta^2 \\ t\zeta^3 & 0 & 0 \end{pmatrix}$$

according to (7). Then [HKL] computed the twisted Alexander polynomial as

$$\Delta_\chi = 4t^2 + (\zeta_5^3 + \zeta_5^2 + 5)t + 4.$$

Note that the two solutions η, λ of Δ_χ are roots of unity $\doteq -0.42274 \pm 0.90624\sqrt{-1}$. Further, they showed that, if K was slice, the Casson-Gordon invariant associated to χ might be zero. Thus, in this situation, the left hand side in Theorem 3.1 should be zero.

To deal with the left hand side, we start by computing $\text{Col}_X^{\text{red}}(D_f)$. Furthermore, we put the assignments of α_1 and α_2 , by setting

$$\mathcal{C}(\alpha_1) = (0, 0, 0, f(\alpha_1)), \quad \mathcal{C}(\alpha_2) = (x, y, z, f(\alpha_2)) \in (\mathbb{C}[t]/\Delta_f)^3 \times G.$$

Denote $\mathcal{C}(\alpha_j)$ by β_j . Then, we can easily see that, these $\mathcal{C}(\alpha_1)$ and $\mathcal{C}(\alpha_2)$ give rise to a coloring over f , if and only if the following holds:

$$\beta_1 \triangleleft \beta_2 = \beta_3, \quad \beta_2 \triangleleft \beta_4 = \beta_5, \quad \beta_4 \triangleleft \beta_1 = \beta_2, \quad \beta_{10} \triangleleft \beta_{12} = \beta_3, \quad \beta_{12} \triangleleft \beta_3 = \beta_{11}, \quad \beta_1 \triangleleft \beta_{11} = \beta_{12},$$

$$\beta_{2i-1} \triangleleft \beta_{2i} = \beta_{2i+1}, \quad \beta_{2i} \triangleleft \beta_{2i-1} = \beta_{2i-2} \quad (3 \leq i \leq 5).$$

Then, by the help of computer program, it can be seen that $\text{Col}_X^{\text{red}}(D_f) \cong \mathbb{C}[t]/\Delta_f$, and that it is generated by such (x, y, z) with

$$y = \frac{8\zeta^2 + \zeta^3 + 8\zeta^4 - 20\zeta^2 t - 20\zeta^4 t}{-12 - \zeta + \zeta^4} x,$$

$$z = \frac{-5 - 24\zeta + 112\zeta^3 + 28\zeta^4 - 60\zeta t + 240\zeta^3 t + 60\zeta^4 t}{4(4 - 13\zeta + 65\zeta^2 + 18\zeta^4)} x.$$

Further, we will compute $\mathcal{Q}_\psi(\mathcal{C}, \mathcal{C}')$ and the associated signature, where ψ is defined by $\psi((x_1, x_2, x_3), (y_1, y_2, y_3)) = \sqrt{-1} \sum x_i \overline{y_i}$. Denote by $\psi(\mathcal{D}_1(\mathcal{C}(\alpha_\tau), \mathcal{D}_2(\mathcal{C}'(\beta_\tau)))$ the term in (7) with $\alpha_\tau = \alpha_i$ and $\beta_\tau = \alpha_j$. Then, $\mathcal{Q}_\psi(\mathcal{C}, \mathcal{C}')$ is, by definition, equal to

$$\Phi(1, 11) + \Phi(12, 3) + \Phi(10, 12) - \Phi(4, 1) - \Phi(1, 2) - \Phi(2, 4) - \left(\sum_{3 \leq i \leq 5} \Phi(2i, 2i-1) + \Phi(2i-1, 2i) \right).$$

However, the resulting computation is a little complicated. Accordingly, to reduce it on the help of computer, it is sensible to consider $H^1(Y; M) \cong \mathbb{C}[t]/(t - \lambda) \oplus \mathbb{C}[t]/(t - \kappa)$ componentwise. Actually, by the help of computer,

$$\mathcal{Q}_\psi(\mathcal{C}, \mathcal{C}')|_{(t-\lambda)} \doteq 29.413653381x\bar{x}', \quad \mathcal{Q}_\psi(\mathcal{C}, \mathcal{C}')|_{(t-\eta)} \doteq -29.413653381x\bar{x}'.$$

As an exercise of linear algebra, we can easily see that the signature of $\text{Tr}(\mathcal{Q}_\psi)$ is equal to the sum of the signatures restricted on the two components. Therefore, by definition, we easily get a table on the signature $\sigma(\lambda(\mathcal{H} \otimes \tau, b \otimes w))$ with respect to $w = \exp(\sqrt{-1}\theta)$ with $0 \leq \theta < 2\pi$, as the table below (Here, the parts denoted by ‘?’ is undefined by non-genericity). In summary, the left hand side mentioned above is not zero. Hence, the assumption of sliceness is false, as desired.

$\exp(\sqrt{-1}\theta)$	1	...	λ	...	κ	...	-1	...	$-\lambda$	...	$-\kappa$	...	1
$\sigma(\lambda(\mathcal{H} \otimes \tau, b \otimes w))$	?	0	?	2	?	0	?	0	?	0	?	0	?

5 Proofs of the statements in Sections 2-3

We will give the proofs of all the statements in Sections 2-3.

5.1 Proofs of the statements in Section 2.2

We will prove Theorems 2.2 and 2.4, by modifying the proofs of [M1, Duality theorem] and [Neu, Theorem in §11]. In this subsection, we suppose the assumption $(\dagger)$ described in §2.2.

First, we will describe strictly the isomorphism Υ_Δ^* stated in Theorem 2.2. Given a CW-complex structure of Y , we have that of $\tilde{Y}$ as a lift. Then, with a choice of $\gamma_0 \in \pi_1(Y)$ representing the covering transformation $\tau : \tilde{Y} \rightarrow \tilde{Y}$, the Reidemeister-Schreier method with respect to $\pi_1(\tilde{Y}) \hookrightarrow \pi_1(Y) \rightarrow \mathbb{Z}$ implies the following identification of cellular complexes

$$V[t^{\pm 1}] \otimes_{f_{\text{pre}}} C_*(Y, \partial Y; \mathbb{Z}) = V \otimes_{f_{\text{pre}}} C_*(\tilde{Y}, \partial \tilde{Y}; \mathbb{Z}).$$

Therefore, we can define a map

$$\Upsilon : C_*(Y, \partial Y; V[t^{\pm 1}]) \longrightarrow C_*(\tilde{Y}, \partial \tilde{Y}; V); \quad \sum_n t^n v_n \otimes c_n \longmapsto \sum_n (v_n \cdot f_{\text{pre}}(\gamma_0^{-n})) \otimes (\tau^n c_n), \quad (10)$$

where $\sum_n t^n v_n$ lies in $V[t^{\pm 1}]$ with $v_n \in V$. This Υ is known to be a chain isomorphism, independent of the choice of γ_0 ; see [KL, Theorem 2.1] or Shapiro’s lemma. Under the assumption $(\dagger)$, consider the module $M := V \otimes_{\mathbb{F}} (\mathbb{F}[t^{\pm 1}]/(\Delta_f))$, and regard it as a local system of $\pi_1(Y)$ via the tensored representation $\alpha \otimes f_{\text{pre}}$. Then, the map Υ subject to the ideal (Δ_f) induces an isomorphism $H_1(Y, \partial Y; M) \xrightarrow{\sim} H_1(\tilde{Y}, \partial \tilde{Y}; V)$ as finite dimensional $\mathbb{F}$ -spaces. Since $\mathbb{F}$ is of characteristic 0, the universal coefficient theorem thus implies that the previous isomorphism in (10) dually induces the isomorphism from $H^1(Y, \partial Y; M)$ to the $\mathbb{Z}$ -equivalent cohomology

$H_{\mathbb{Z}}^1(\tilde{Y}, \partial\tilde{Y}; M)$. Since the image is naturally isomorphic to $H^1(\tilde{Y}, \partial\tilde{Y}; V)$ by the trace map Tr , we consequently have

$$\Upsilon_{\Delta}^* : H^1(Y, \partial Y; M) \xrightarrow{\sim} H^1(\tilde{Y}, \partial\tilde{Y}; V).$$

Further, when $M = \mathbb{F}[t]/\Delta_f$, we can construct a homomorphism $\Upsilon_{\Delta}^* : H^2(Y, \partial Y; \mathbb{F}[t]/\Delta_f) \rightarrow H^2(\tilde{Y}, \partial\tilde{Y}; \mathbb{F})$.

Next, let us observe two filtrations of $\tilde{Y}$ such as [M1, Neu, Ka]. Choose a circle valued Morse function $\bar{\alpha} : Y \rightarrow S^1$, which represents $\alpha : \pi_1(Y) \rightarrow \mathbb{Z}$. Then, we can lift it to $\tilde{\alpha} : \tilde{Y} \rightarrow \mathbb{R}$, and define a surface $\Sigma \subset \tilde{Y}$ to be the preimage of a regular value $p_0 \in \mathbb{R}$, that is, $\Sigma := \tilde{\alpha}^{-1}(p_0)$. Then, for $q \in \mathbb{Z}$, the two filtrations are defined as

$$N_q^+ := \tilde{\alpha}^{-1}((-\infty, q + p_0)), \quad \text{and} \quad N_q^- := \tilde{\alpha}^{-1}((-q + p_0, \infty)).$$

Consider the Mayer-Vietoris sequence of the triple $(\tilde{Y}, \partial\tilde{Y} \cup N_q^+, \partial\tilde{Y} \cup N_q^-)$ written in

$$\cdots \rightarrow H^i(\tilde{Y}, \partial\tilde{Y}; V) \xrightarrow{\delta_c^*} H^{i+1}(\tilde{Y}, \partial\tilde{Y} \cup N_q^+ \cup N_q^-; V) \rightarrow \bigoplus H^{i+1}(\tilde{Y}, \partial\tilde{Y} \cup N_q^{\pm}; V) \rightarrow \cdots \quad (11)$$

We further discuss the middle term and δ_c . Notice that, by definition, $\tilde{Y} \setminus (N_p^+ \cup N_q^-)$ is compact and $\cap_{p \in \mathbb{Z}} N_p^+ = \cap_{q \in \mathbb{Z}} N_q^- = \emptyset$. Since V is over a field $\mathbb{F}$ of characteristic 0, the inductive limit of the second term as $q \rightarrow -\infty$ can be shown to be explicitly the relative cohomology of $\tilde{Y}$ with compact support (see [FS] for the details on $\lim^1$). Further, denoting by j the inclusion $\Sigma \subset \tilde{Y}$, consider the composite

$$H^i(\tilde{Y}, \partial\tilde{Y}; V) \xrightarrow{j^*} H^i(\Sigma, \partial\Sigma; V) \xrightarrow{j_!} H_{\text{compact}}^{i+1}(\tilde{Y}, \partial\tilde{Y}; V), \quad (12)$$

where the second map is the Gysin map (or pushforward). Similar to [Neu, §12], we can easily see, in comparison with the Thom class of a normal bundle along Σ , that the map δ_c^* is equal to (12), up to isomorphisms. Precisely, $a_i \circ \delta_c^* = j_! \circ j^*$ for some automorphism a_i on $H_{\text{compact}}^{i+1}(\tilde{Y}, \partial\tilde{Y}; V)$. Hence, as the projection formula, we obtain the multiplicative identity

$$a_{\deg(x \smile y)} \circ \delta_c^*(x \smile y) = j_!(j^*x \smile j^*y) = (j_! \circ j^*(x)) \smile y = (a_{\deg(x)} \circ \delta_c^*(x)) \smile y. \quad (13)$$

Further, we will examine this δ_c^* in an algebraic way. For this, let $\mathbb{F}[[\mathbb{Z}]]$ be the $\mathbb{F}[t^{\pm 1}]$ -module consisting of Laurent series $\sum_{k=-\infty}^{\infty} b_k t^k$ with $b_k \in \mathbb{F}$. Further, recall from §2.2 the Laurent polynomial rings $\mathbb{F}[[\mathbb{Z}]]_{\pm}$ and the embeddings $\iota_{\pm} : \mathbb{F}((t)) \rightarrow \mathbb{F}[[\mathbb{Z}]]_{\pm}$. Then, the map $\iota_+ - \iota_-$ defines $q : \mathbb{F}((t))/\mathbb{F}[t^{\pm 1}] \rightarrow \mathbb{F}[[\mathbb{Z}]]$ by passage. Since $\mathbb{F}[[\mathbb{Z}]]_+ \cap \mathbb{F}[[\mathbb{Z}]]_- = \mathbb{F}[t^{\pm 1}]$, we thus have a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{F}[t^{\pm 1}] & \xrightarrow{\quad} & \mathbb{F}((t)) & \xrightarrow{\quad} & \mathbb{F}((t))/\mathbb{F}[t^{\pm 1}] \longrightarrow 0 \\ & & \parallel & & \downarrow \iota_+ \oplus \iota_- & & \downarrow q \\ 0 & \longrightarrow & \mathbb{F}[t^{\pm 1}] & \xrightarrow{\mathfrak{s}} & \mathbb{F}[[\mathbb{Z}]]_+ \oplus \mathbb{F}[[\mathbb{Z}]]_- & \xrightarrow{\mathfrak{t}} & \mathbb{F}[[\mathbb{Z}]] \longrightarrow 0 \end{array} \quad \begin{array}{l} \text{(exact)} \\ \\ \text{(exact),} \end{array} \quad (14)$$

where $\mathfrak{s}$ and $\mathfrak{t}$ are defined by the maps $x \mapsto (x, x)$ and $(x, y) \mapsto x - y$, respectively. Let δ_*^u and δ_*^b denote by the respective connecting maps obtained from the upper and bottom sequences. Notice that, similar to [Neu, Page 170] or [FS], the chain isomorphism Υ in (10) gives rise to an isomorphism

$$\Upsilon_*^{\text{comp}} : H_j(Y, \partial Y; V[[\mathbb{Z}]]) \xrightarrow{\sim} H_j^{\text{compact}}(\tilde{Y}, \partial\tilde{Y}; V)$$

with compact support. Moreover, the connecting maps δ_*^b and δ_*^c satisfy $\Upsilon_* \circ \delta_*^b = \delta_*^c \circ \Upsilon_*^{\text{comp}}$, by construction.

We shall make a further examination in the case where V is the trivial coefficient $\mathbb{Q}$, in more details. It was shown [M1, §4] that the inverse limit $\lim_{q \rightarrow -\infty} H_i(\tilde{Y}, \partial \tilde{Y} \cup N_q^\pm; \mathbb{Q})$ for any i vanishes. Therefore, dually, the connecting map $\delta_*^c : H_{i+1}^{\text{comp}}(\tilde{Y}, \partial \tilde{Y}; \mathbb{Q}) \rightarrow H_i(\tilde{Y}, \partial \tilde{Y}; \mathbb{Q})$ is an isomorphism; By (12), the fundamental class $\tilde{\mu}_3$ in the third homology $H_3^{\text{comp}}(\tilde{Y}, \partial \tilde{Y}; \mathbb{Q}) \cong \mathbb{Q}$ is sent to the orientation class $[\Sigma] \in H_2(\tilde{Y}, \partial \tilde{Y}; \mathbb{Q})$ such that $(p_\alpha)_*[\Sigma] = \mu_\alpha$ and $[\Sigma] = \tau_*[\Sigma]$ (see also [Ka, Lemmas 6.1–6.4] as the integral case). Furthermore, by duality, let us compute this second homology $H_2(Y, \partial Y; \mathbb{Q}[t^{\pm 1}])$ as

$$H^1(Y; \mathbb{Q}[t^{\pm 1}]) \cong \text{Hom}_{\mathbb{Q}[t^{\pm 1}]\text{-mod}}(H_1(Y; \mathbb{Q}[t^{\pm 1}]), \mathbb{Q}[t^{\pm 1}]) \oplus \text{Ext}^1(H_0(Y; \mathbb{Q}[t^{\pm 1}]), \mathbb{Q}[t^{\pm 1}]).$$

Notice that the last term is $\mathbb{Q}[t^{\pm 1}]/(t-1)$; denote the generator by $\tilde{\mu}_\alpha^{\mathbb{Q}[t^{\pm 1}]}$, which satisfies $\mu_\alpha = \tilde{\mu}_\alpha^{\mathbb{Q}[t^{\pm 1}]}|_{t^0}$. Moreover, since $H_*(Y, \partial Y; \mathbb{Q}((t)))$ is a free $\mathbb{Q}((t))$ -module by definition, the long exact sequence obtained from (14) ensures some $\mu_3^{\text{fund}} \in H_3(Y, \partial Y; \mathbb{Q}((t))/\mathbb{Q}[t^{\pm 1}])$ satisfying

$$\tilde{\mu}_\alpha^{\mathbb{Q}[t^{\pm 1}]} = \delta_*^u(\mu_3^{\text{fund}}), \quad \Upsilon_*^{\text{comp}} \circ q_*(\mu_3^{\text{fund}}) = \tilde{\mu}_3.$$

To summarize, we can compute $\delta_*^c(\tilde{\mu}_3)$ as

$$\delta_*^c(\tilde{\mu}_3) = \delta_*^c \circ \Upsilon_*^{\text{comp}} \circ q_*(\mu_3^{\text{fund}}) = \Upsilon_* \circ \delta_*^b \circ q_*(\mu_3^{\text{fund}}) = \Upsilon_* \circ \delta_*^u(\mu_3^{\text{fund}}) = \Upsilon_*(\tilde{\mu}_\alpha^{\mathbb{Q}[t^{\pm 1}]}). \quad (15)$$

Using the above observation, we now be in a situation to prove Theorem 2.2.

Proof of Theorem 2.2. The finiteness of the dimension of $H^1(Y, \partial Y; M)$ can be easily verified by using that of $H_1(Y, \partial Y; M)$ and the universal coefficients theorem.

Let us show the equality $\text{Tr} \circ \mathcal{Q}_\psi = \smile_{KL} \circ \Upsilon_\Delta^{\otimes 2}$ in the statement. By (15) and the definition of $\smile_{KL}$, for $x, y \in H^1(Y, \partial Y; M)$, we can compute $\Upsilon_\Delta^*(x) \smile_{KL} \Upsilon_\Delta^*(y)$ as

$$\begin{aligned} \psi_{\text{pre}}(\Upsilon_\Delta^*(x \smile y) \frown [\Sigma]) &= \psi_{\text{pre}}(\Upsilon_\Delta^*(x \smile y) \frown \delta_*^c(\tilde{\mu}_3)) \\ &= \psi_{\text{pre}}(\Upsilon_\Delta^*(x \smile y) \frown \Upsilon_*(\tilde{\mu}_\alpha^{\mathbb{Q}[t^{\pm 1}]}) \\ &= \psi_{\text{pre}}(q_*((x \smile y) \frown \tilde{\mu}_\alpha^{\mathbb{Q}[t^{\pm 1}]})|_{t^0}) \end{aligned}$$

From the definition of the trace map and q_* , the last term is $\text{Tr} \circ \mathcal{Q}_\psi(x, y)$, as desired. $\square$

Next, we will prove Theorem 2.4 by showing the following lemma:

Lemma 5.1 (cf. [M1, Assumption 8]). *Let $\mathbb{F}$ be a field with $\text{Char} \mathbb{F} = 0$. Assume that $H_1(Y; V[t^{\pm 1}])$ is of finite dimension. Then, the above connecting map with $i = 1$*

$$\delta_c^* : H^1(\tilde{Y}, \partial \tilde{Y}; V) \longrightarrow H_{\text{compact}}^2(\tilde{Y}, \partial \tilde{Y}; V)$$

is injective.

Proof. Notice that every element ν in $H_1(\tilde{Y}, \partial \tilde{Y}; V)$ is represented by a 1-cycle contained in $N_{q_{\pm, \nu}}^\pm$ for some $q_{\pm, \nu} \in \mathbb{N}$. Since the homology $H_1(\tilde{Y}, \partial \tilde{Y}; V)$ is of finite dimension by assumption, the maximal s of such $q_{j, \nu}$'s admits the surjectivity of the natural map

$$H_1(N_{-s}, N_0 \cup (\partial \tilde{Y} \cup N_0); V) \longrightarrow H_1(\tilde{Y}, \partial \tilde{Y} \cup N_0; V).$$

Therefore, by observing some long exact sequence, the natural map

$$H_1(\tilde{Y}, \partial\tilde{Y} \cup N_0; V) \longrightarrow H_1(\tilde{Y}, \partial\tilde{Y} \cup N_{-s}; V)$$

is zero; therefore we have $\lim_{q \rightarrow -\infty} H_1(\tilde{Y}, \partial\tilde{Y} \cup N_q^\pm; V) \cong 0$. Thus, the dual vanishing $\lim_{q \rightarrow -\infty} H^1(\tilde{Y}, \partial\tilde{Y} \cup N_q^\pm; V) \cong 0$ follows from the universal coefficient theorem and $\text{Char}\mathbb{F} = 0$. Hence, we obtain the desired injectivity from the above Mayer-Vietoris sequence (11). $\square$

Proof of Theorem 2.4. Recall from Remark 2.3 the equality $[i_+^{-1}(\sum_{j \geq 0} \Upsilon_\Delta^*(\tau_*^{-j}x) \smile_{KL} \Upsilon_\Delta^*(y) \cdot t^j)] = [\mathcal{Q}_\psi(x, y)/\Delta_f]$, and the injectivity of i_+ . Therefore, it is sufficient for the proof to show the desired non-degeneracy of $\mathcal{Q}_\psi$.

Next, we will show (16) below. Set the natural maps $\iota : (Y, \emptyset) \rightarrow (Y, \partial Y)$ and $\tilde{\iota} : (\tilde{Y}, \emptyset) \rightarrow (\tilde{Y}, \partial\tilde{Y})$. Recall from Assumption $(\dagger)$ that $H^1(Y, \partial Y; M)$ is of finite dimension over $\mathbb{F}$ with $\text{Char}\mathbb{F} = 0$. Thus, the universal coefficient theorem gives an isomorphism between finite dimensional $\mathbb{F}$ -spaces

$$\text{Ker}(\iota^* : H^1(Y, \partial Y; M) \rightarrow H^1(Y; M)) \cong \text{Ker}(\iota_* : H_1(Y; V[t^{\pm 1}]) \rightarrow H_1(Y, \partial Y; V[t^{\pm 1}])).$$

Further, via the chain isomorphisms Υ_* and Υ_Δ^* , this isomorphism can be written in

$$\text{Ker}(\tilde{\iota}^* : H^1(\tilde{Y}, \partial\tilde{Y}; V) \rightarrow H^1(\tilde{Y}; V)) \cong \text{Ker}(\tilde{\iota}_* : H_1(\tilde{Y}; V) \rightarrow H_1(\tilde{Y}, \partial\tilde{Y}; V)). \quad (16)$$

Consequently, the left kernel is a dual module of the right one by the Kronecker product.

Using the previous duality in (16), let us show the non-degeneracy of the bilinear form $\smile_{KL}$ on the kernel $\text{Ker}(\tilde{\iota}^*)$. For $x, y \in H^1(\tilde{Y}, \partial\tilde{Y}; V)$, compute $\psi_{\text{pre}}\langle \tilde{\iota}^*(x) \smile y, [\Sigma] \rangle \in \mathbb{F}$ as

$$\begin{aligned} \psi_{\text{pre}}\langle a_2 \circ a_1^{-1} \circ \delta_c^*(\tilde{\iota}^*(x) \smile y), \tilde{\mu}_3 \rangle &= \psi_{\text{pre}}\langle \delta_c^*(\tilde{\iota}^*(x)) \smile y, \tilde{\mu}_3 \rangle \\ &= \psi_{\text{pre}}\langle y, \tilde{\mu}_3 \smile \delta_c^*(\tilde{\iota}^*(x)) \rangle, \end{aligned}$$

where we use (13). Note that $\tilde{\mu}_3 \smile \delta_c^*$ is an injection by Poincaré-duality and Lemma 5.1. From the definition of $\smile_{KL}$ and the duality in (16), it is non-degenerate on the kernel $\text{Ker}(\iota^*)$, as desired. $\square$

5.2 Proofs of Theorem 2.1

Changing the subject, the end of this section is devoted to proving Theorem 2.1 which recovers the Blanchfield pairing from $\mathcal{Q}_\psi$. Let Λ denote the ring $\mathbb{Z}[t^{\pm 1}]$, and Y_K denote the knot complement $S^3 \setminus K$ in what follows.

For the purpose, let us briefly review from [T2] a matrix presentation of the pairing. For a knot K with $\#K = 1$, we choose a Seifert surface, that is, a connected compact surface Σ embedded in S^3 such that the boundary is K . Take the intersection form $I_\Sigma : H_1(\Sigma; \mathbb{Z})^{\otimes 2} \rightarrow \mathbb{Z}$. For $u \in H_1(\Sigma; \mathbb{Z})$, let $\iota_+(u)$ be the resulting 1-cycle of pushing u a small distance along the positive normal to Σ . Then, we define a bilinear form $\theta : H_1(\Sigma; \mathbb{Z})^{\otimes 2} \rightarrow \mathbb{Z}$ by setting $\theta(u \otimes v) = I_\Sigma(\iota_+(u), v) \in \mathbb{Z}$. With a choice of a basis $v_1, \dots, v_{2g} \in H_1(\Sigma; \mathbb{Z})$, the integral matrix V defined by $\{\theta(v_i, v_j)\}_{i,j \leq 2g}$ is referred to as a *Seifert matrix* of Σ . Then, it is well-known that the Alexander module $H_1(S^3 \setminus K; \Lambda)$ is finitely presented by the matrix $tV - V'$

with coefficient Λ , i.e.,

$$\Lambda^{2g} \xrightarrow{tV-V'} \Lambda^{2g} \longrightarrow H_1(S^3 \setminus K; \Lambda) \longrightarrow 0 \quad (\text{exact}), \quad (17)$$

where V' is the transpose of V , and that the Alexander polynomial $\Delta_K \in \mathbb{Z}[t^{\pm 1}]$ is equal to $\det(tV - V')$ up to multiplication by $\pm t^{\pm 1}$. Further, it is shown [T2, Proposition 2.5] that the Blanchfield pairing is given by a bilinear form $H_1(Y_K; \Lambda)^{\otimes 2} \rightarrow \Lambda/(\Delta_K)$ with the matrix $(1-t)\text{adj}(t^{-1}V - V') \cdot (\Delta_K^{-1} \cdot \text{id}_{\Lambda^{2g}})$.

Further, let us translate the Blanchfield pairing from the viewpoints of first cohomologies, following [T1]. Let $\Delta : \Lambda \rightarrow \Lambda$ be multiplication by Δ_K . Since $H^1(Y_K, \partial Y_K; \Lambda) \cong H_2(Y_K; \Lambda) \cong 0$ is known, the short exact sequence of coefficient groups $0 \rightarrow \Lambda \xrightarrow{\Delta} \Lambda \rightarrow \Lambda/(\Delta_K) \rightarrow 0$ derives a part of the long exact sequence

$$0 \longrightarrow H^1(Y_K, \partial Y_K; \Lambda/(\Delta_K)) \xrightarrow{\beta^*} H^2(Y_K, \partial Y_K; \Lambda) \xrightarrow{\Delta^*} H^2(Y_K, \partial Y_K; \Lambda). \quad (18)$$

Here, the Cayley-Hamilton theorem implies that Δ^* annihilates $H^2(Y_K, \partial Y_K; \Lambda)$; hence, the Bockstein map β^* is an isomorphism onto. Following the Poincaré duality $\mathcal{P} : H^2(Y_K, \partial Y_K; \Lambda) \cong H_1(Y_K; \Lambda)$ and (17), every element of $H^2(Y_K, \partial Y_K; \Lambda)$ is represented as someone in Λ^{2g} . Then it is shown [T1, p.487] that, for any representative $\vec{b} \in \Lambda^{2g}$, the image $(\beta^*)^{-1}(\vec{b})$ is in 1:1-correspondence with $\text{adj}(t^{-1}V - V')\vec{b}'$ in the sense of a 1-cochain with coefficients $\Lambda/(\Delta_K)$ which represents an element of $H^1(Y_K, \partial Y_K; \Lambda/(\Delta_K))$. Hence, thanks to the composite isomorphism

$$(\mathcal{P})^{-1} \circ (\beta^*)^{-1} : H^1(Y_K, \partial Y_K; \Lambda/(\Delta_K)) \cong H_1(Y_K; \Lambda), \quad (19)$$

the Blanchfield pairing valued in $\Lambda/(\Delta_K)$ on the first cohomology $H^1(Y_K, \partial Y_K; \Lambda/(\Delta_K))$ is given by the matrix

$$(1-t^{-1}) \cdot (tV - V') \quad \text{modulo } \Delta_K. \quad (20)$$

Proof of Theorem 2.1. First, the former statement $H^1(Y_K, \partial Y_K; \Lambda) \cong H_1(Y_K; \Lambda)$ readily follows from the isomorphism β^* in (18) and the Poincaré duality.

To complete the proof, we will compute $\psi_0 \langle x \smile y, [\Sigma] \rangle \in \Lambda/\Delta_K$ in more details, where ψ_0 is $(\Lambda/\Delta_K)^2 \rightarrow \Lambda/\Delta_K$ which sends (a, b) to $\bar{a}b$. Recall from (15) the 2-class $\mu_\alpha^{\mathbb{Q}[t^{\pm 1}]} \in H_2(Y, \partial Y; \mathbb{Q}[t^{\pm 1}])$ satisfying $\delta_*^u(\mu_\alpha^{\mathbb{Q}[t^{\pm 1}]}) = \mu_3^{\text{fund}}$ and $\mu_\alpha^{\mathbb{Q}[t^{\pm 1}]}|_{t^0} = [\Sigma] = \mu_\alpha$. Furthermore, notice the Leibniz rule of the Bockstein map:

$$\beta^*(x \smile y) = \beta^*(x) \smile y - x \smile \beta^*(y) \quad \text{for any } x, y \in H^1(Y_K, \partial Y_K; \Lambda/(\Delta_K)).$$

Thus, via the inclusion $\Lambda/(\Delta_K) \hookrightarrow \mathbb{Q}((t))/\mathbb{Q}[t^{\pm 1}]$, we compute $\psi_0 \langle x \smile y, [\Sigma] \rangle$ as

$$\psi_0 \langle x \smile y, \mu_\alpha \rangle = \psi_0 \langle x \smile y, \delta_*^u(\mu_\alpha^{\mathbb{Q}[t^{\pm 1}]}) \rangle = \tilde{\psi}_1 \langle \beta^*(x) \smile y, \mu_3^{\text{fund}} \rangle - \tilde{\psi}_2 \langle x \smile \beta^*(y), \mu_3^{\text{fund}} \rangle,$$

where we define $\tilde{\psi}_1 : \Lambda \otimes \Lambda/(\Delta_K) \rightarrow \Lambda/(\Delta_K)$ and $\tilde{\psi}_2 : \Lambda/(\Delta_K) \otimes \Lambda \rightarrow \Lambda/(\Delta_K)$ as the canonical lifts of ψ_0 . Furthermore, the cup product between these H^1 and H^2 is shown to be the canonical inner product (see [T1, Proposition 4.1] as the viewpoint of Alexander-Whitney

map). Hence, by the above discussion on β^* , this pairing $\psi_0\langle x \smile y, [\Sigma] \rangle$ has the following matrix presentation:

$$\left(\overline{x \cdot (t^{-1}V - V')} \right) \cdot y' - \bar{x} \cdot (y \cdot (t^{-1}V - V'))',$$

which turns out to be $(1+t)\bar{x}(V - t^{-1}V')y'$. In comparison with (20), we have the conclusion $\mathcal{Q}_{\psi_0}(x, y) = \psi_0\langle x \smile y, [\Sigma] \rangle = (1-t)^{-1}(1+t) \cdot \text{Bl}_K(x, y)$ as required. $\square$

5.3 Proof of Theorem 3.1

This section completes the proof of Theorem 3.1. As seen below, the proof is essentially based on some results in [Neu] and [KL].

Proof of Theorem 3.1. Let Y_∞ be the infinite cover of X_q associated with the free abelianization $\alpha : \pi_1(X_q) \rightarrow \text{Free}H_1(X_q; \mathbb{Z}) \cong \mathbb{Z}$, and $\mu_\alpha \in H_2(Y_\infty; \mathbb{Z})$ be a lift of the dual element of $[\alpha] \in H^1(Y_\infty; \mathbb{Z})$. Let $\psi_{\text{pre}} : \mathbb{C}^2 \rightarrow \mathbb{C}$ send (x, y) to $\bar{x}y$. Then, in analogous to (3), we can consider the cohomology $H_t^1(Y_\infty; \mathbb{C})$ twisted by $\chi \otimes \hat{\alpha}_w$, and set up the 2-form

$$\smile_{\psi_{\text{pre}}} : H_t^1(Y_\infty; \mathbb{C})^{\otimes 2} \xrightarrow{\sim} H_t^2(Y_\infty; \mathbb{C} \otimes \mathbb{C}) \xrightarrow{\langle \psi_{\text{pre}}, \bullet \rangle} H_t^2(Y_\infty; \mathbb{C}) \xrightarrow{\cap \mu_\alpha} \mathbb{C}. \quad (21)$$

Then, by Lemma 5.2 below, this 2-form is equal to $\text{Tr} \circ \mathcal{Q}_{\psi_q}$ on the twisted cohomology $\mathcal{H}$.

Since the representations χ and $\hat{\alpha}_w$ of $\pi_1(X_q)$ are unitary, this situation on $\text{Tr} \circ \mathcal{Q}_{\psi_q}$ satisfies the conditions in Theorem 3.2 in [Neu] (Here, we apply $\alpha = \hat{\alpha}_w$ and $\beta = \hat{\chi}$). Hence, the desired equality is directly obtained from the theorem. $\square$

Furthermore, we will show the proposition used above:

Lemma 5.2. *Let us use the notation above, and suppose the assumptions in Theorem 3.1.*

Then, the skew hermitian form $\text{Tr} \circ \mathcal{Q}_\psi$ on the twisted cohomology $\mathcal{H} = H_t^1(S^3 \setminus K, \partial S^3 \setminus K; M)$ over $\mathbb{C}$ is isomorphic to $\smile_{\psi_{\text{pre}}}$ on $H^1(Y_\infty; \mathbb{C}_\chi)$.

Proof. We will get the desired isomorphism by constructing three isomorphisms. Denote by M_χ the extended system $\mathbb{C}_\chi[t]/(\Delta_\chi)$ of E_q or X_q which comes from the character χ and $\bar{\alpha} : \pi_1(X_q) \rightarrow \pi_1(E_q) \rightarrow \mathbb{Z}$. Since f_χ is the induced representation from χ , the Shapiro Lemma (see [HKL, §3]) directly yields the canonical ring isomorphism

$$p^* : H_t^*(S^3 \setminus K, \partial S^3 \setminus K; M_\Delta) \xrightarrow{\sim} H_t^*(E_q, \partial E_q; M_\chi).$$

Next, notice that the longitude $\mathbf{l}_K \in \pi'_K$ is bounded by a lift of a surface μ_α . Therefore, this $\mathbf{l}_K$ vanishes in $\pi'_K/\pi''_K = H_1(\pi_K; \mathbb{Z}[t^{\pm 1}])$; Thus, $f_\chi(\mathbf{l}_K) = \text{id}$ by definition. Then, thank to the excision axiom, the inclusion $i : E_q \hookrightarrow X_q$ gives rise to the second isomorphism

$$i^* : H_t^*(X_q, X_q \setminus \text{Int}(E_q); M_\chi) \xrightarrow{\sim} H_t^*(E_q, \partial E_q; M_\chi).$$

Finally, we will construct the third one. Denote the space $X_q \setminus \text{Int}(E_q)$ by cX_q for short. Consider the natural map $\eta : (X_q, \emptyset) \rightarrow (X_q, cX_q)$, which induces the sequence

$$\cdots \rightarrow H_t^0(cX_q) \rightarrow H_t^1(X_q, cX_q) \xrightarrow{\eta^*} H_t^1(X_q) \rightarrow H_t^1(cX_q) \rightarrow \cdots \quad (\text{exact}),$$

in coefficients M_χ . We now compute the both edges. Notice that cX_q is homotopic to S^1 as a meridian $\tilde{\mathbf{m}}$, and that $f_\chi(\tilde{\mathbf{m}}) = t \in M_\chi$ by definition. Therefore, by assumption (vi), the zero-th $H_t^0(cX_q; M_\chi)$ is the invariant part $M_\chi^\mathbb{Z} = 0$, and $H_t^1(cX_q; M_\chi)$ is the coinvariant $M_\chi/(t-1) \cong \mathbb{C}[t]/(t-1)$. In conclusion, the above sequence is rewritten in

$$0 \longrightarrow H_t^1(X_q, cX_q) \xrightarrow{\eta^*} H_t^1(X_q) \longrightarrow \mathbb{C}[t]/(t-1) \longrightarrow 0 \quad (\text{split exact}).$$

In particular, the cup product in the $(t-1)$ -part of $H_t^*(X_q)$ is trivial. Hence, we have the third isomorphism $H_t^1(X_q, cX_q) \xrightarrow{\eta^*} H_t^1(X_q)/(1-t)$ modulo $(1-t)$.

In summary, the composite $\eta^* \circ (i^*)^{-1} \circ p^*$ means $H_t^1(S^3 \setminus K, \partial S^3 \setminus K; M_\Delta) \cong H_t^1(X_q; M_\chi)$. Further, the above isomorphisms preserve the ring structures. Hence, Theorem 2.2 immediately concludes the desired equality $\text{Tr} \circ \mathcal{Q}_\psi = \smile_{\psi_{\text{pre}}}$. $\square$

A Appendix: comparison with the work of Goldman

Starting from the setting in Goldman's symplectic structure [Go], we will geometrically observe (Proposition A.1) the associated 2-form $\mathcal{Q}_\psi$ (1) from the viewpoint of flat bundles. Here are the settings: Let G be a semi-simple reductive Lie group over $\mathbb{R}$ or $\mathbb{C}$, and let the Lie algebra $\mathfrak{g}$ be with the adjoint action of G . Take the Killing form $\psi_{\text{pre}} : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathbb{C}$ as a non-degenerate symmetric form.

Next, we will define a subset (22) of homomorphisms $\pi_1(Y) \rightarrow G$, for some compact 3-manifold Y . Choose a surjection $\alpha : \pi_1(Y) \rightarrow \mathbb{Z}$, and take the associated infinite cyclic covering $p_\alpha : \tilde{Y} \rightarrow Y$. Then, for an $f_{\text{pre}} : \pi_1(Y) \rightarrow G$, we obtain a representation $f : \pi_1(Y) \rightarrow \text{End}(\mathfrak{g}[t^{\pm 1}])$ via $f_{\text{pre}} \otimes \alpha$; As in §2.2, we choose a twisted Alexander polynomial $\Delta_f \in \mathbb{C}[t^{\pm 1}]$, and denote the discriminant by $\text{disc}(\Delta_f) \in \mathbb{C}$. Further, consider $\text{dis}_{Y,G} := \sup_{f_{\text{pre}}} (\dim H^1(\tilde{Y}; \mathfrak{g})) \in \mathbb{N} \cup \{\infty\}$, where f_{pre} runs over all homomorphisms $\pi_1(Y) \rightarrow G$. Let us define the subset of $\text{Hom}(\pi_1(Y), G)$ to be

$$\left\{ f_{\text{pre}} \in \text{Hom}(\pi_1(Y), G) \left| \begin{array}{l} \text{dis}_{Y,G} = \dim H^1(\tilde{Y}; \mathfrak{g}), \quad \text{disc}(\Delta_f) \neq 0 \quad \text{and} \\ \mathfrak{g}(1 - f_{\text{pre}}(g)) = \mathfrak{g}, \quad \text{for any } g \in \pi_1(\partial Y) \end{array} \right. \right\} \quad (22)$$

and denote it by $R_G(Y)$. We can easily see that $\det(\text{id}_{\mathfrak{g}} - f_{\text{pre}}(g))$ is an algebraic function on $\text{Hom}(\pi_1(Y), G)$. Hence, $R_G(Y)$ is Zariski open dense in $\text{Hom}(\pi_1(Y), G)$, possibly $R_G(Y) = \emptyset$.

To proceed the discussion, let us assume the finiteness $\text{dis}_{Y,G} < \infty$; we can obtain such many examples from the fibered α , that is, a homeomorphism $\tilde{Y} \cong \Sigma_\alpha \times \mathbb{R}$ for some compact surface Σ_α . Under the assumption, for $f_{\text{pre}} \in R_G(Y)$, we have isomorphisms

$$H^1(\tilde{Y}; \mathfrak{g}) \cong H^1(\tilde{Y}, \partial \tilde{Y}; \mathfrak{g}) \cong H^1(Y, \partial Y; \mathfrak{g}[t]/(\Delta_f)),$$

where the first one is obtained from the property $\mathfrak{g}(1 - f_{\text{pre}}(g)) = \mathfrak{g}$, and the second follows from Theorems 2.2. Moreover, Theorem 2.4 implies that this space has a non-degenerate skew-hermitian 2-form defined as $\smile_{KL}$. Here recall the well-known fact (see [Go]) that the cohomology $H^1(\tilde{Y}; \mathfrak{g})$ is naturally identified with the tangent space of $\text{Hom}(\pi_1(\tilde{Y}), G)/G$ at $f_{\text{pre}} \circ p_\alpha$. In other ward, the 2-form is regarded as a symplectic structure on the flat moduli

space $\text{Hom}(\pi_1(\tilde{Y}), G) // G$ away from appreciate singular points. Furthermore, consider the continuous map

$$p_\alpha^* : \text{Hom}(\pi_1(Y), G) / G \longrightarrow \text{Hom}(\pi_1(\tilde{Y}), G) / G$$

induced from the covering $p_\alpha : \tilde{Y} \rightarrow Y$. Then, we can easily construct a vector bundle on the above space $R_G(Y)$ whose fiber is $H^1(\tilde{Y}; \mathfrak{g})$, and denote it by E .

From the viewpoint of the base space Y , it follows from Theorem 2.2 that the 2-form $\smile_{KL}$ is described as $\text{Tr} \circ \mathcal{Q}_\psi$ via p_α^* . To summarize, we can state the desired geometric meaning:

Proposition A.1. *Assume $\text{dist}_{Y,G} < \infty$ as above. Let $E \rightarrow R_G(Y)$ be the above vector bundle constructed with fiber $H^1(\tilde{Y}; \mathfrak{g})$.*

Then, the non-degenerate skew-hermitian 2-form $\text{Tr} \circ \mathcal{Q}_\psi$ on E is equal to the pullback via p_α^* of the symplectic structure on the flat moduli space $\text{Hom}(\pi_1(\tilde{Y}), G) // G$ from appreciate singular points.

To conclude, let us sum up as follows. Some previous papers (see, e.g., [Gu]) dealt with $H^1(Y; \mathfrak{g})$ in coefficients $\mathfrak{g}$, as a Lagrangian subspace of the Goldman symplectic structure on $H^1(\partial Y; \mathfrak{g})$ from the boundaries surfaces ∂Y . In particular, the 2-form on $\pi_1(Y) \rightarrow G$ is often zero. However, concerning Proposition A.1, as seen in the proof of Theorem 2.2, if Y is $S^3 \setminus L$, the 2-form in Proposition A.1 arises from the symplectic structure on $H^1(\Sigma; \mathfrak{g})$, where Σ is a Seifert surface. Furthermore, Theorem 4.1 makes the non-degenerate 2-form $\text{Tr} \circ \mathcal{Q}_\psi$ computable. Hence, we hope that this work will be of use in analysing the non-zero 2-form on the flat moduli space $\text{Hom}(\pi_1(\tilde{Y}), G) // G$ on the infinite cover $\tilde{Y}$.

B Universality of Alexander quandle cocycle invariants.

The paper [CJKLS] constructed a knot invariant from quandle cocycles. However, the invariant was defined in a combinatorial way without topological meanings. This appendix reviews the invariant, and gives a topological meaning in the Alexander case, as an application of Theorem 2.1.

For the purpose, we first briefly review the invariant, assuming knowledges in §4.1. Denote the abelianization $\pi_1(S^3 \setminus K) \rightarrow \mathbb{Z}$ of a knot K by f , let X be a $\mathbb{Z}[t^{\pm 1}]$ -module. Then, the pair of X and the binary operation $(x, y) \mapsto tx + y - ty$ is called the Alexander quandle; see [CJKLS]. Further, we suppose a map $\phi : X^2 \rightarrow A$ satisfying the equality

$$\phi(x, z) - \phi(y, z) - \phi(tx - ty + y, z) + \phi(tx - tz + z, ty - tz + z) = 0, \quad (23)$$

for any $x, y, z \in X$. Such a map ϕ is called a *quandle 2-cocycle* [CJKLS]. Then, in analogy of §4.1, we can take a coloring set $\text{Col}_X(D_f)$, and define a map

$$\mathcal{I}_\phi : \text{Col}_X(D_f) \longrightarrow A, \quad \mathcal{C} \longmapsto \sum_{\tau} \epsilon_{\tau} \cdot \phi(\mathcal{C}(\alpha_{\tau}), \mathcal{C}(\beta_{\tau})),$$

where τ ranges over all the crossings of D , and the symbols $\alpha_{\tau}, \beta_{\tau}$ are the arcs and $\epsilon_{\tau} \in \{\pm 1\}$ is the sign of τ according to Figure 1. Then, it is known [CJKLS] that, thanks to (23), the map $\mathcal{I}_\phi$ is independent of the choice of D by (23); then, $\mathcal{I}_\phi$ is called *the quandle cocycle invariant*.

For example, given an additive homomorphism $\psi : X^2 \rightarrow A$ satisfying (23), we can easily verify, by definitions, that the map ϕ_ψ defined by $\phi_\psi(x, y) = \psi(x - y, y - yt^{-1})$ satisfies (23), and that the invariant $\mathcal{I}_\Phi$ is equal to the restricted 2-form $\mathcal{Q}_\psi \circ \Delta$, where Δ is the diagonal map $\text{Col}_X(D_f) \rightarrow \text{Col}_X(D_f)^2$.

Furthermore, we review the S -equivalences of knots. Two knots K and K' are S -equivalent² if they are related by a finite sequence of the (double delta) local move shown in Figure 4. Furthermore, Trotter [T2] classically showed that K and K' are S -equivalent if and only if the associated Blanchfield pairings are isomorphic as bilinear forms.




Figure 4: A double delta move with 24 arcs

Theorem B.1. *Two knots K and K' are S -equivalent, if and only if for every Alexander quandle X and every quandle 2-cocycle $\phi : X^2 \rightarrow A$, the associated cocycle invariants $\mathcal{I}_\Phi$ and $\mathcal{I}'_\Phi$ are equivalent.*

In conclusion, this theorem from the viewpoint of the Blanchfield pairing concisely concludes a topological meaning in the Alexander case.

Proof. Thanks to Proposition B.2 below, if two knots K and K' are S -equivalent, then the resulting cocycle invariants are equal.

Conversely, we assume the equivalence of the resulting cocycle invariants with $X = A = \mathbb{Z}[t^{\pm 1}]/\Delta_K$ and ψ_0 as stated in Theorem 2.1. Since $t + 1$ is divisible by the nonzero integer $\Delta_K(-1)$ in X , the assumption and Theorem 2.1 imply the equivalence of the two Blanchfield pairings. Hence, by the result of Trotter [T2] above, K and K' are S -equivalent, as desired. $\square$

Proposition B.2. (i) *For any bilinear function $\psi : X^2 \rightarrow A$ satisfying $\psi(x, y) = \psi(ty, x)$, the associated bilinear form $\mathcal{Q}_\psi$ is an S -equivalent knot invariant.*

(ii) *For any quandle 2-cocycle $\phi : X^2 \rightarrow A$, the cocycle invariant is also S -equivalent.*

Proof. It is enough for (i) to show the local invariance with respect to the double delta move. Given an X -coloring of D , consider X -colors of the arcs α_i with $1 \leq i \leq 12$ in Figure 4. By direct calculation, it can be seen that the correspondence $\alpha_i \leftrightarrow \beta_i$ ensures another coloring of D' , which yields a $\mathbb{Z}[t^{\pm 1}]$ -module isomorphism $\mathcal{B} : \text{Col}_X(D_f) \cong \text{Col}_X(D'_f)$. Moreover, together with Lemma B.3 and the assumption $\psi(a, b) = \psi(tb, a)$, one easily verifies that the desired equality $\mathcal{Q}_\psi(\mathcal{C}_1, \mathcal{C}_2) = \mathcal{Q}'_\psi(\mathcal{B}(\mathcal{C}_1), \mathcal{B}(\mathcal{C}_2)) \in A$ holds for any X -colorings $\mathcal{C}_1$ and $\mathcal{C}_2$.

(ii) According to [Joy, Theorem 17.3] concerning “the Abelianization of the knot quandle”, for any Alexander quandle X , there is a functorial $\mathbb{Z}[t^{\pm 1}]$ -module isomorphism

$$\text{Col}_X(D_f) \cong \text{Hom}_{\mathbb{Z}[t^{\pm 1}]\text{-mod}}(H_1(S^3 \setminus K; \mathbb{Z}[t^{\pm 1}]), X) \oplus X.$$

²As a result of [NS], the definition is equal to the common one by means of Seifert matrices up to some transformations; see also [T2].

In other ward, $\text{Col}_X(D_f)$ is representable by the Alexander module $H_1(S^3 \setminus K; \mathbb{Z}[t^{\pm 1}])$. Hence, by the universality, we may assume $X = H_1(S^3 \setminus K; \mathbb{Z}[t^{\pm 1}])$ hereafter. In particular, $(1-t)X = X$.


Further, consider “the quandle second homology $H_2^Q(X; \mathbb{Z})$ ” defined in [CJKLS], which is isomorphic to the quotient $\mathbb{Z}$ -module

$$G_X := X \otimes_{\mathbb{Z}} X / \{x \otimes y - (ty) \otimes x \mid x, y \in X\}.$$

This result is essentially due to Clauwens [Cla] (see also [No1, §5] for the proof). Further, consider a homomorphism $\psi_{\text{uni}} : X \otimes X \rightarrow G_X$ which sends (x, y) to $[x \otimes y]$. Then, it is known (see, e.g., [CJKLS, No1]), every cocycle invariant $\Delta \circ \mathcal{Q}_\psi$ factors through the homology $H_2^Q(X; \mathbb{Z})$. Precisely, there is an additive homomorphism $P_\psi : G_X \rightarrow A$ such that $\mathcal{Q}_{\psi_{\text{uni}}} \circ \Delta = P_\psi(\mathcal{Q}_\psi \circ \Delta)$ for any link L . Hence, from the definitions of G_X and ψ_{uni} , the proof for the S -equivalence is done in the same manner as (i). $\square$

Since the following lemmas is elementary, we omit writing details of the proofs.

Lemma B.3. *Given an X -coloring of the eight arcs illustrated in the figure below, the sum of weights with respect to the four crossings is $\psi((1-t)(a-b), c-d) = \psi((1-t)(u-v), d-c) \in A$.*



Acknowledgments

The author sincerely expresses his gratitude to Takahiro Kitayama and Masaaki Wada for valuable comments. The work is partially supported by JSPS KAKENHI Grant Number 00646903.

References

- [AS] M. F. Atiyah, I.M.Singer, *The index of elliptic operators: III*, Ann. of Math, **87** (1968), 546–604.
- [Bla] R. Blanchfield, *Intersection theory of manifolds with operators with applications to knot theory*, Ann. of Math. **65**: (1957) 340–356
- [Cla] F.J.-B.J. Clauwens, *The adjoint group of an Alexander quandle*, arXiv:math/1011.1587.
- [CG] A. J. Casson, C. McA. Gordon, *Cobordism of classical knots*, mimeographed notes (1975)
- [CJKLS] J. S. Carter, D. Jelsovsky, S. Kamada, L. Langford, M. Saito, *Quandle cohomology and state-sum invariants of knotted curves and surfaces*, Trans. Amer. Math. Soc. **355** (2003), 3947–3989.
- [COT] T. D. Cochran, K. Orr, P. Teichner, *Knot concordance, whitney towers and L^2 -signatures*, Ann. of Math. **157** (2003) 433–519.
- [CHL] T. D. Cochran, S. Harvey, C. Leidy, *Knot concordance and higher-order Blanchfield duality*, Geom. Topol. **13** (2009), 1419–1482.
- [FG] V. Florens, P. Gilmer, *On the slice genus of links*, Algebr. Geom. Topol. **3** (2003), 905–920.

- [Gi] P. Gilmer, *On the slice genus of knots*, Invent. Math. **66** (1982), 191–197.
- [FKK] S. Friedl, T. Kim, T. Kitayama, *Poincaré duality and degrees of twisted Alexander polynomials*, Indiana Univ. Math. J. **61** (2012), 147–192.
- [FS] M. Farber and D. Schütz, *Moving Homology Classes to Infinity*, Forum Math. **19** (2007), no. 2, 281–296.
- [Gu] S. Gukov, *Three-Dimensional Quantum Gravity, Chern-Simons Theory, and the A-Polynomial*, Commun. Math. Phys. **255** (2005), 577–627.
- [Go] W. Goldman, *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*, Invent. math. **85** (1986) 263–302.
- [Gi] P. Gilmer, *On the slice genus of knots*, Invent. Math. **66** (1982), 191–197.
- [Hil] J. Hillman, *Algebraic invariants of links*, second edition, Series on Knots and everything. **32** World Scientific (2002).
- [HKL] C. Herald, P. Kirk and C. Livingston, *Metabelian representations, twisted Alexander polynomials, knot slicing, and mutation*, Preprint (2008), to appear in Math. Z.
- [HSW] J. Hillman, D. Silver, S. Williams, *On reciprocity of twisted Alexander invariants*, Algebr. Geom. Topol. **10** (2010), 1017–1026.
- [Joy] D. Joyce, *A classifying invariant of knots, the knot quandle*, J. Pure Appl. Algebra **23** (1982), 37–65.
- [Ka] A. Kawauchi, *Three dualities on the integral homology of infinite cyclic coverings of manifolds*, Osaka J. Math. **23** (1986), 633–651.
- [KL] P. Kirk, C. Livingston. *Twisted Alexander invariants, Reidemeister torsion, and Casson-Gordon invariants*, Topology, **38** (1999), 635–661.
- [Lin] X.S. Lin, *Representations of knot groups and twisted Alexander polynomials*, Acta Math. Sin. (Engl. Ser.) **17** (2001), 361–380.
- [NS] S. Naik, T. Stanford, *A move on diagrams that generates S -equivalence of knots*, J. Knot Theory Ramifications **12** (2003), 717–724.
- [Neu] W. Neumann, *Signature related invariants of manifolds-I. Monodromy and γ -invariants*, Topology **18** (1979), 147–172.
- [No1] T. Nosaka, *Homotopical interpretation of link invariants from finite quandles*, preprint.
- [No2] ———, *Twisted cohomology pairings of knots I; diagrammatic computation*, preprint.
- [Mat] T. Matumoto, *On the signature invariants of a non-singular complex sesquilinear form*, J. Math. Soc. Japan **29** (1977), no. 1, 67–71.
- [M1] J. W. Milnor, *Infinite cyclic coverings*, Conference on the Topology of Manifolds (Michigan State Univ., E. Lansing, Mich., 1967), Prindle, Weber & Schmidt, Boston, Mass., (1968) 115–133.
- [M2] ———, *On isometries of inner product spaces*, Invent. Math. **8**, (1969) 83–97.
- [T1] H. F. Trotter, *Homology of group systems with applications to knot theory*, Ann. of Math. **76** (1962), 464–498.
- [T2] ———, *On S -equivalence of Seifert matrices*, Invent. Math. **20** (1973), 173–207.
- [W] M. Wada, *Twisted Alexander polynomial for finitely presentable groups*, Topology **33** (1994), 241–256.

Faculty of Mathematics, Kyushu University, 744, Motooka, Nishi-ku, Fukuoka, 819-0395, Japan

E-mail address: nosaka@math.kyushu-u.ac.jp